



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:38 PM UTC

PDB ID : 4UTR / pdb_00004utr
Title : Crystal structure of zebrafish Sirtuin 5 in complex with glutarylated CPS1-peptide
Authors : Pannek, M.; Gertz, M.; Steegborn, C.
Deposited on : 2014-07-22
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	 4% 76% 17% • 5%
1	B	275	 % 78% 13% •• 7%
2	C	9	 11% 56% 44%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-DEPENDENT PROTEIN DEACYLASE SIRTUIN-5, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	262	Total	C	N	O	S	0	2	0
			2051	1293	371	372	15			
1	B	257	Total	C	N	O	S	0	1	0
			2003	1265	361	362	15			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	GLY	-	expression tag	UNP Q6DHI5
A	25	ILE	-	expression tag	UNP Q6DHI5
A	26	ASP	-	expression tag	UNP Q6DHI5
A	27	PRO	-	expression tag	UNP Q6DHI5
A	28	PHE	-	expression tag	UNP Q6DHI5
A	29	THR	-	expression tag	UNP Q6DHI5
B	24	GLY	-	expression tag	UNP Q6DHI5
B	25	ILE	-	expression tag	UNP Q6DHI5
B	26	ASP	-	expression tag	UNP Q6DHI5
B	27	PRO	-	expression tag	UNP Q6DHI5
B	28	PHE	-	expression tag	UNP Q6DHI5
B	29	THR	-	expression tag	UNP Q6DHI5

- Molecule 2 is a protein called GLUTARYL-CPS1 PEPTIDE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	9	Total	C	N	O	0	0	0
			68	47	9	12			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	BEZ	-	modified residue	UNP Q5R209

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 6 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Na	0	0
			1	1		

- Molecule 7 is GLUTARIC ACID (CCD ID: GUA) (formula: C₅H₈O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			8	5	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	39	Total	O	0	0
			39	39		
8	B	18	Total	O	0	0
			18	18		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: NAD-DEPENDENT PROTEIN DEACYLASE SIRTUIN-5, MITOCHONDRIAL



• Molecule 1: NAD-DEPENDENT PROTEIN DEACYLASE SIRTUIN-5, MITOCHONDRIAL



• Molecule 2: GLUTARYL-CPS1 PEPTIDE



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	87.22Å 87.22Å 314.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	75.53 – 2.90 75.53 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (75.53-2.90) 99.9 (75.53-2.90)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.213 , 0.262 0.216 , 0.256	Depositor DCC
R_{free} test set	830 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	58.2	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4206	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GUA, BEZ, EDO, NA, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	0/2106	1.05	6/2855 (0.2%)
1	B	0.90	1/2059 (0.0%)	1.09	8/2789 (0.3%)
2	C	0.92	0/60	1.14	1/79 (1.3%)
All	All	0.90	1/4225 (0.0%)	1.07	15/5723 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	292	THR	C-O	5.19	1.30	1.24

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	65	THR	N-CA-C	-8.82	102.10	113.12
1	A	63	VAL	CB-CA-C	-8.45	101.46	111.18
2	C	4	LYS	CG-CD-CE	-7.06	95.06	111.30
1	A	65	THR	N-CA-C	-6.92	104.48	113.12
1	A	206	LYS	N-CA-C	-6.59	97.09	108.56
1	B	225	SER	N-CA-C	6.26	118.11	111.28
1	B	273	GLU	N-CA-C	6.13	122.81	113.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	209	ASN	N-CA-C	6.01	120.20	112.86
1	A	209	ASN	N-CA-C	5.78	119.92	112.86
1	A	225	SER	N-CA-C	5.58	117.37	111.28
1	B	63	VAL	CB-CA-C	-5.58	104.46	111.04
1	B	205	ARG	CA-C-O	-5.36	114.28	120.49
1	B	272	MET	N-CA-C	-5.23	99.66	110.80
1	A	104	VAL	N-CA-CB	5.08	119.34	110.65
1	B	104	VAL	N-CA-CB	5.04	119.27	110.65

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	278	THR	Peptide
1	B	70	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2051	0	2007	26	0
1	B	2003	0	1968	22	0
2	C	68	0	66	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	8	0	12	0	0
4	B	4	0	6	0	0
5	B	4	0	6	0	0
6	B	1	0	0	0	0
7	C	8	0	0	0	0
8	A	39	0	0	1	0
8	B	18	0	0	0	0
All	All	4206	0	4065	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:VAL:HG12	1:A:149[B]:HIS:CD2	2.21	0.75
1:A:235:LEU:HD13	1:A:256:PHE:HB3	1.72	0.71
1:B:100:TYR:O	1:B:104:VAL:HG13	1.96	0.66
1:A:100:TYR:O	1:A:104:VAL:HG13	1.98	0.64
1:B:111:ASN:H	1:B:114:HIS:HD2	1.44	0.64
1:A:54:GLY:HA2	1:A:136:GLN:HG2	1.83	0.60
1:A:233:ARG:HD3	8:A:2037:HOH:O	2.01	0.60
1:A:115:LEU:HD11	1:B:115:LEU:HD11	1.82	0.60
1:A:71:GLY:O	1:A:77:GLN:HG2	2.03	0.57
1:A:165:CYS:SG	1:A:205:ARG:NH1	2.77	0.57
1:A:103:GLU:OE1	1:A:106:ARG:NH1	2.41	0.53
1:B:65:THR:O	1:B:66:PHE:HB2	2.08	0.53
1:B:111:ASN:H	1:B:114:HIS:CD2	2.26	0.53
1:A:111:ASN:HB2	1:A:112:PRO:CD	2.39	0.52
1:A:251:TYR:CD2	1:A:252:PRO:HA	2.45	0.52
1:A:40:ARG:NH2	1:A:298:GLU:OXT	2.42	0.52
1:A:251:TYR:C	1:A:251:TYR:CD1	2.88	0.51
1:A:132:VAL:HG12	1:A:149[B]:HIS:CG	2.46	0.51
1:A:65:THR:O	1:A:66:PHE:HB2	2.11	0.50
1:B:251:TYR:CD1	1:B:252:PRO:HA	2.47	0.50
1:B:111:ASN:HB2	1:B:112:PRO:CD	2.41	0.50
1:A:272:MET:HE3	1:A:287:GLY:HA2	1.95	0.49
1:A:235:LEU:CD1	1:A:256:PHE:HB3	2.42	0.48
1:B:54:GLY:HA2	1:B:136:GLN:HG2	1.95	0.48
1:A:185:ALA:HB3	1:A:190:THR:HG21	1.94	0.48
1:B:105:MET:HE2	1:B:139:ASP:HB2	1.95	0.47
1:A:111:ASN:HB2	1:A:112:PRO:HD2	1.95	0.47
1:A:105:MET:HE2	1:A:139:ASP:HB2	1.97	0.47
1:B:273:GLU:O	1:B:274:CYS:HB2	2.13	0.47
1:A:246:THR:OG1	1:A:247:SER:N	2.47	0.47
1:A:109:MET:HE2	1:B:116:ALA:N	2.30	0.47
1:B:185:ALA:HB3	1:B:190:THR:HG21	1.97	0.46
1:B:239:ASP:OD1	1:B:239:ASP:N	2.47	0.46
1:B:233:ARG:O	1:B:236:GLU:HG2	2.15	0.45
1:B:246:THR:OG1	1:B:247:SER:N	2.49	0.45
1:B:133:ILE:HD12	1:B:133:ILE:N	2.32	0.44
1:A:109:MET:HE2	1:B:115:LEU:C	2.42	0.44
1:B:111:ASN:HB2	1:B:112:PRO:HD2	1.99	0.44
1:A:251:TYR:HB2	2:C:6:TYR:CE2	2.53	0.43
1:A:133:ILE:HD12	1:A:133:ILE:N	2.33	0.43
1:B:48[B]:HIS:HB3	1:B:238:CYS:HA	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:ASN:OD1	1:A:284:HIS:HE1	2.02	0.42
1:B:161:ARG:HH11	1:B:215:HIS:CE1	2.38	0.41
1:B:65:THR:O	1:B:66:PHE:CB	2.68	0.41
1:B:48[A]:HIS:HB2	1:B:238:CYS:HA	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/275 (94%)	250 (96%)	8 (3%)	2 (1%)	16	44
1	B	254/275 (92%)	246 (97%)	8 (3%)	0	100	100
2	C	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
All	All	521/559 (93%)	502 (96%)	17 (3%)	2 (0%)	30	59

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	207	SER
1	A	276	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/229 (96%)	203 (93%)	16 (7%)	13	38
1	B	214/229 (93%)	203 (95%)	11 (5%)	21	53
2	C	6/6 (100%)	4 (67%)	2 (33%)	0	0
All	All	439/464 (95%)	410 (93%)	29 (7%)	15	43

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	104	VAL
1	A	136	GLN
1	A	189	ASN
1	A	205	ARG
1	A	207	SER
1	A	216	VAL
1	A	223	LEU
1	A	238	CYS
1	A	250	VAL
1	A	259	GLN
1	A	272	MET
1	A	273	GLU
1	A	275	THR
1	A	278	THR
1	A	282	LYS
1	A	289	CYS
1	B	37	THR
1	B	77	GLN
1	B	79	GLN
1	B	104	VAL
1	B	130	SER
1	B	136	GLN
1	B	205	ARG
1	B	238	CYS
1	B	250	VAL
1	B	273	GLU
1	B	289	CYS
2	C	3	LEU
2	C	8	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	136	GLN
1	A	284	HIS
1	B	99	HIS
1	B	114	HIS
1	B	284	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	EDO	A	1301	-	3,3,3	0.55	0	2,2,2	0.57	0
4	EDO	A	1302	-	3,3,3	0.26	0	2,2,2	0.83	0
5	DMS	B	1301	-	3,3,3	0.56	0	3,3,3	1.06	0
4	EDO	B	1300	-	3,3,3	0.54	0	2,2,2	0.23	0
7	GUA	C	1299	2	6,7,8	1.60	1 (16%)	6,7,9	1.14	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	1301	-	-	1/1/1/1	-
4	EDO	A	1302	-	-	1/1/1/1	-
7	GUA	C	1299	2	-	2/5/5/6	-
4	EDO	B	1300	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	1299	GUA	C2-C1	2.61	1.56	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	1299	GUA	C3-C2-C1	-2.42	108.18	114.51

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	1300	EDO	O1-C1-C2-O2
4	A	1302	EDO	O1-C1-C2-O2
7	C	1299	GUA	O2-C1-C2-C3
4	A	1301	EDO	O1-C1-C2-O2
7	C	1299	GUA	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/275 (95%)	0.02	11 (4%) 40 32	16, 52, 89, 126	2 (0%)
1	B	257/275 (93%)	0.19	3 (1%) 76 69	29, 60, 105, 126	1 (0%)
2	C	8/9 (88%)	0.02	1 (12%) 8 7	44, 52, 67, 74	0
All	All	527/559 (94%)	0.10	15 (2%) 55 46	16, 55, 99, 126	3 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	256	PHE	4.0
1	A	282	LYS	3.3
1	A	277	ALA	3.3
1	B	71	GLY	3.0
1	A	275	THR	2.7
1	A	69	PRO	2.7
1	A	258	PRO	2.5
1	B	75	LYS	2.5
2	C	8	VAL	2.5
1	A	71	GLY	2.5
1	A	278	THR	2.4
1	A	257	ALA	2.1
1	A	149[A]	HIS	2.1
1	B	206	LYS	2.0
1	A	259	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EDO	A	1301	4/4	0.82	0.20	52,55,56,60	0
4	EDO	B	1300	4/4	0.86	0.16	54,54,59,67	0
7	GUA	C	1299	8/9	0.86	0.16	44,53,65,67	0
6	NA	B	1302	1/1	0.88	0.20	60,60,60,60	1
4	EDO	A	1302	4/4	0.91	0.20	56,57,57,57	0
5	DMS	B	1301	4/4	0.95	0.12	47,49,52,54	0
3	ZN	B	1299	1/1	0.97	0.06	86,86,86,86	0
3	ZN	A	1300	1/1	1.00	0.03	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.