



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:49 PM UTC

PDB ID : 5V6Z / pdb_00005v6z
Title : Crystal structure of human caspase-7 soaked with allosteric inhibitor 2-{[2-(4-chlorophenyl)-2-oxoethyl]sulfanyl}benzoic acid
Authors : Vance, N.R.; Gakhar, L.; Spies, M.A.
Deposited on : 2017-03-17
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	12% 57% 9% 34%
1	B	303	8% 56% 13% 31%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3175 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caspase-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	201	Total	C	N	O	S	0	0	0
			1478	936	253	275	14			
1	B	210	Total	C	N	O	S	0	0	0
			1605	1021	272	297	15			

- Molecule 2 is 2-{[2-(4-chlorophenyl)-2-oxoethyl]sulfanyl}benzoic acid (CCD ID: 8YJ) (formula: C₁₅H₁₁ClO₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	Cl	O	S	0	0
			20	15	1	3	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	20	Total	O	0	0
			20	20		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	52	Total	O	0	0
			52	52		

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.34Å 89.34Å 184.90Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	59.34 – 2.60 59.34 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (59.34-2.60) 99.9 (59.34-2.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575)	Depositor
R, R_{free}	0.204 , 0.249 0.210 , 0.253	Depositor DCC
R_{free} test set	1418 reflections (5.25%)	wwPDB-VP
Wilson B-factor (Å ²)	64.1	Xtriage
Anisotropy	0.515	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3175	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 8YJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/1499	0.82	0/2019
1	B	0.56	0/1632	0.85	1/2199 (0.0%)
All	All	0.54	0/3131	0.84	1/4218 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	294	MET	CA-CB-CG	5.43	124.96	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1478	0	1357	19	0
1	B	1605	0	1517	25	0
2	B	20	0	0	2	0
3	A	20	0	0	0	0
3	B	52	0	0	2	0
All	All	3175	0	2874	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:294:MET:HE3	2:B:401:8YJ:CAC	2.19	0.73
1:A:292:VAL:HG12	1:A:294:MET:HE1	1.73	0.70
1:B:233:ARG:HA	1:B:239:SER:HA	1.75	0.69
1:B:246:CYS:O	1:B:250:GLU:HG3	1.98	0.64
1:A:292:VAL:HG12	1:A:294:MET:CE	2.30	0.61
1:A:123:LEU:HD12	1:A:166:PHE:CE1	2.36	0.60
1:A:123:LEU:HD12	1:A:166:PHE:HE1	1.67	0.59
1:B:294:MET:HE3	2:B:401:8YJ:CAD	2.33	0.58
1:B:169:ASP:OD1	1:B:170:ARG:HG3	2.04	0.57
1:A:60:TYR:OH	1:A:218:ASP:OD1	2.24	0.56
1:B:93:ASP:HA	3:B:501:HOH:O	2.03	0.56
1:B:97:LEU:HD13	1:B:140:ILE:HG21	1.87	0.55
1:B:163:THR:HG21	1:B:221:PHE:HE2	1.74	0.52
1:A:111:TYR:CG	1:A:122:LEU:HD21	2.44	0.52
1:B:54:ARG:HB3	1:B:59:GLN:HE22	1.74	0.52
1:A:163:THR:HG21	1:A:221:PHE:HE2	1.74	0.52
1:A:158:PRO:HG2	1:A:161:ASP:OD2	2.10	0.51
1:B:84:MET:HB3	1:B:144:HIS:CD2	2.46	0.51
1:A:114:CYS:SG	1:A:118:LYS:HE2	2.52	0.50
1:B:84:MET:HE2	1:B:144:HIS:CG	2.47	0.49
1:A:244:ALA:O	1:A:248:ILE:HG12	2.13	0.49
1:B:124:LYS:O	1:B:128:GLU:HG3	2.13	0.49
1:B:243:GLN:HA	1:B:243:GLN:OE1	2.13	0.48
1:B:266:ASN:OD1	1:B:289:PRO:HB2	2.13	0.48
1:A:177:LYS:O	1:A:179:LYS:HE3	2.14	0.47
1:B:184:GLN:HA	1:B:224:SER:HB3	1.96	0.47
1:B:115:SER:O	1:B:119:MET:HG3	2.15	0.47
1:B:76:LYS:HB2	1:B:90:THR:HG21	1.97	0.46
1:B:112:ASN:HB3	3:B:528:HOH:O	2.14	0.46
1:B:96:ALA:O	1:B:100:CYS:HB2	2.15	0.46
1:B:60:TYR:CD1	1:B:178:PRO:HD3	2.51	0.46
1:B:146:GLU:O	1:B:147:GLU:C	2.57	0.46
1:A:267:ASP:HB2	1:B:296:THR:O	2.16	0.45
1:A:65:GLU:OE1	1:A:133:ASN:ND2	2.49	0.45
1:B:256:LEU:HD23	1:B:256:LEU:HA	1.87	0.44
1:B:98:PHE:CD1	1:B:98:PHE:C	2.95	0.44
1:A:267:ASP:OD1	1:A:271:ARG:NH2	2.51	0.43
1:A:73:ILE:HB	1:A:141:LEU:HD23	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:TYR:CD1	1:A:178:PRO:HD3	2.54	0.42
1:A:178:PRO:HG3	1:A:299:LEU:HD13	2.01	0.41
1:A:184:GLN:HA	1:A:224:SER:HB3	2.03	0.41
1:A:271:ARG:HH22	1:B:297:LYS:NZ	2.18	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/303 (64%)	189 (98%)	4 (2%)	0	100	100
1	B	202/303 (67%)	199 (98%)	3 (2%)	0	100	100
All	All	395/606 (65%)	388 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/267 (53%)	142 (100%)	0	100	100
1	B	164/267 (61%)	161 (98%)	3 (2%)	51	76
All	All	306/534 (57%)	303 (99%)	3 (1%)	68	86

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	98	PHE
1	B	108	VAL
1	B	156	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	ASN
1	A	120	GLN
1	A	184	GLN
1	B	88	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	8YJ	B	401	-	21,21,21	0.92	0	28,28,28	1.57	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8YJ	B	401	-	-	12/13/13/13	0/2/2/2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	8YJ	CAH-SAG-CAF	-5.83	92.29	102.27
2	B	401	8YJ	CAA-CAF-SAG	3.41	124.34	120.36
2	B	401	8YJ	CAE-CAF-SAG	-2.47	115.51	121.48
2	B	401	8YJ	CAI-CAH-SAG	-2.11	104.44	111.36

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	8YJ	SAG-CAH-CAI-CAJ
2	B	401	8YJ	CAH-CAI-CAJ-CAK
2	B	401	8YJ	CAH-CAI-CAJ-CAO
2	B	401	8YJ	OAS-CAI-CAJ-CAK
2	B	401	8YJ	OAS-CAI-CAJ-CAO
2	B	401	8YJ	SAG-CAH-CAI-OAS
2	B	401	8YJ	CAF-CAA-CAP-OAQ
2	B	401	8YJ	CAF-CAA-CAP-OAR
2	B	401	8YJ	CAA-CAF-SAG-CAH
2	B	401	8YJ	CAB-CAA-CAP-OAQ
2	B	401	8YJ	CAB-CAA-CAP-OAR
2	B	401	8YJ	CAI-CAH-SAG-CAF


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	8YJ	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	201/303 (66%)	0.86	35 (17%) 4 3	49, 72, 109, 127	4 (1%)
1	B	210/303 (69%)	0.32	23 (10%) 10 8	41, 60, 91, 108	5 (2%)
All	All	411/606 (67%)	0.58	58 (14%) 6 5	41, 64, 106, 127	9 (2%)

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	272	HIS	9.9
1	A	234	SER	9.5
1	B	53	ASP	6.0
1	A	235	PRO	5.6
1	B	234	SER	5.4
1	A	79	ASP	5.2
1	A	147	GLU	5.1
1	A	77	ASN	5.1
1	A	84	MET	5.0
1	A	88	ASN	4.8
1	A	232	TRP	4.8
1	B	56	PRO	4.8
1	A	273	PHE	4.5
1	A	272	HIS	4.4
1	A	186	CYS	4.4
1	A	78	PHE	4.0
1	A	82	THR	4.0
1	B	233	ARG	4.0
1	A	215	VAL	3.9
1	A	287	GLN	3.8
1	A	231	SER	3.8
1	A	152	GLY	3.8
1	B	286	LYS	3.7
1	A	233	ARG	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	226	VAL	3.6
1	A	150	ILE	3.5
1	B	225	THR	3.5
1	A	57	THR	3.4
1	B	187	ARG	3.3
1	A	153	LYS	3.3
1	B	232	TRP	3.3
1	A	81	VAL	3.2
1	A	151	TYR	3.0
1	A	236	GLY	2.9
1	A	157	THR	2.9
1	B	273	PHE	2.9
1	A	148	ASN	2.8
1	B	231	SER	2.8
1	B	215	VAL	2.8
1	B	237	ARG	2.8
1	B	51	THR	2.7
1	A	185	ALA	2.7
1	B	235	PRO	2.7
1	A	223	TYR	2.7
1	B	55	VAL	2.5
1	A	224	SER	2.4
1	A	80	LYS	2.3
1	B	240	TRP	2.3
1	A	63	ASN	2.3
1	A	237	ARG	2.3
1	B	214	PRO	2.2
1	A	90	THR	2.2
1	B	54	ARG	2.2
1	B	148	ASN	2.1
1	B	236	GLY	2.1
1	A	83	GLY	2.1
1	B	223	TYR	2.1
1	A	156	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	8YJ	B	401	20/20	0.78	0.33	66,82,90,107	20

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 8YJ B 401:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.