



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:09 PM UTC

PDB ID : 7W7Y / pdb_00007w7y
Title : The crystal structure of human abl1 kinase domain in complex with ABL2-A5
Authors : Zhu, C.; Zhang, Z.M.
Deposited on : 2021-12-06
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 5% 82% 12% . .
1	B	276	 8% 82% 12% . .

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4741 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase ABL1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	269	Total	C	N	O	P	S	0	0	0
			2159	1390	351	400	1	17			
1	B	267	Total	C	N	O	P	S	0	0	0
			2138	1378	345	398	1	16			

- Molecule 2 is 5-[3-(5-methanoyl-2-methoxy-4-oxidanyl-phenyl)-1 {H}-pyrrolo[2,3-b]pyridin-5-yl]- {N}, {N}-dimethyl-pyridine-3-carboxamide (CCD ID: 8IW) (formula: C₂₃H₂₀N₄O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			30	23	4	3		
2	B	1	Total	C	N	O	0	0
			30	23	4	3		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	225	Total 225	O 225	0	0
3	B	159	Total 159	O 159	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Tyrosine-protein kinase ABL1

Chain A: 



• Molecule 1: Tyrosine-protein kinase ABL1

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	56.52Å 105.35Å 132.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.97 – 2.20 42.97 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.97-2.20) 99.9 (42.97-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.31 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.9_1692+SVN	Depositor
R, R_{free}	0.175 , 0.201 (Not available) , 0.192	Depositor DCC
R_{free} test set	2054 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	28.8	Xtriage
Anisotropy	0.562	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4741	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 8IW, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2197	0.82	3/2977 (0.1%)
1	B	0.45	0/2174	0.77	3/2945 (0.1%)
All	All	0.48	0/4371	0.80	6/5922 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	357	LYS	N-CA-C	10.40	126.12	113.23
1	B	253	TYR	N-CA-C	6.91	122.81	113.97
1	A	438	SER	CA-C-N	5.60	125.55	119.78
1	A	438	SER	C-N-CA	5.60	125.55	119.78
1	B	397	ALA	CA-C-N	5.36	131.34	121.70
1	B	397	ALA	C-N-CA	5.36	131.34	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	380	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2159	0	2075	26	0
1	B	2138	0	2056	27	0
2	A	30	0	0	1	0
2	B	30	0	0	0	0
3	A	225	0	0	5	2
3	B	159	0	0	7	0
All	All	4741	0	4131	51	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:351:MET:SD	3:A:884:HOH:O	2.18	1.00
1:B:393:PTR:O1P	3:B:701:HOH:O	1.99	0.80
1:A:249:GLY:O	3:A:701:HOH:O	2.00	0.78
1:A:357:LYS:O	3:A:702:HOH:O	2.01	0.78
1:B:397:ALA:HB3	1:B:398:GLY:HA3	1.71	0.72
1:B:386:ARG:NH1	3:B:703:HOH:O	2.23	0.71
1:B:397:ALA:CB	1:B:398:GLY:HA3	2.23	0.68
1:A:237:MET:HE3	1:A:303:GLY:HA3	1.75	0.67
1:B:237:MET:HE3	1:B:303:GLY:HA3	1.83	0.60
1:A:373:GLU:OE2	3:A:703:HOH:O	2.15	0.60
1:B:257:TYR:OH	3:B:702:HOH:O	2.17	0.59
1:A:279:GLU:HG2	1:B:342:TYR:OH	2.02	0.59
1:A:304:VAL:HG13	1:A:306:THR:HG23	1.85	0.57
1:A:351:MET:HE3	1:A:379:VAL:HG21	1.86	0.56
1:B:253:TYR:H	1:B:254:GLY:HA2	1.71	0.55
1:B:253:TYR:N	1:B:254:GLY:HA2	2.21	0.55
1:B:254:GLY:HA2	3:B:754:HOH:O	2.06	0.55
1:B:292:GLU:CD	1:B:386:ARG:HH22	2.15	0.54
1:B:304:VAL:HG13	1:B:306:THR:HG23	1.89	0.53
1:A:493:PHE:HA	1:A:496:MET:HE3	1.94	0.50
1:A:282:GLU:CD	3:A:711:HOH:O	2.55	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:GLU:OE1	1:A:306:THR:N	2.37	0.48
1:A:243:THR:OG1	1:A:262:LYS:HD2	2.14	0.48
1:A:290:MET:HE3	1:A:381:ASP:HB3	1.95	0.48
1:A:319:THR:HG21	1:A:373:GLU:OE2	2.13	0.48
1:B:305:CYS:HB2	1:B:312:TYR:HB2	1.95	0.47
1:B:358:ASN:HA	3:B:762:HOH:O	2.15	0.47
1:A:249:GLY:HA3	1:A:251:GLY:N	2.30	0.45
1:A:271:LYS:NZ	2:A:601:8IW:OBE	2.50	0.45
1:A:249:GLY:HA3	1:A:251:GLY:H	1.82	0.45
1:B:460:ARG:NH2	3:B:707:HOH:O	2.32	0.44
1:B:309:PRO:HA	1:B:310:PRO:C	2.43	0.44
1:A:276:ASP:HA	1:A:277:THR:C	2.43	0.44
1:A:356:LYS:HB2	1:A:356:LYS:HE2	1.74	0.44
1:B:271:LYS:HB3	1:B:313:ILE:CG2	2.48	0.43
1:B:248:LEU:HD12	1:B:248:LEU:HA	1.79	0.43
1:A:295:HIS:CG	1:A:296:PRO:HD2	2.54	0.43
1:A:358:ASN:OD1	1:A:358:ASN:N	2.52	0.43
1:B:248:LEU:HD11	1:B:317:PHE:HE1	1.83	0.43
1:A:354:LEU:O	1:A:358:ASN:HA	2.19	0.42
1:A:438:SER:HA	1:A:439:PRO:HD3	1.83	0.42
1:B:357:LYS:HB3	1:B:357:LYS:HE3	1.88	0.42
1:A:263:LYS:HD2	1:A:264:TYR:CE2	2.54	0.42
1:B:233:ASP:OD1	1:B:234:LYS:N	2.51	0.41
1:B:315:THR:HG21	1:B:382:PHE:HE1	1.85	0.41
1:B:391:ASP:HB2	3:B:722:HOH:O	2.21	0.41
1:B:232:TYR:HA	1:B:235:TRP:HB2	2.02	0.41
1:B:238:GLU:O	1:B:241:ASP:HB2	2.21	0.41
1:A:238:GLU:H	1:A:238:GLU:CD	2.19	0.40
1:B:438:SER:HA	1:B:439:PRO:HD3	1.95	0.40
1:A:386:ARG:HD2	1:B:500:SER:C	2.46	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:822:HOH:O	3:A:822:HOH:O[2_565]	1.51	0.69
3:A:794:HOH:O	3:A:847:HOH:O[2_565]	2.08	0.12

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/276 (96%)	256 (96%)	9 (3%)	1 (0%)	30	34
1	B	262/276 (95%)	253 (97%)	8 (3%)	1 (0%)	30	34
All	All	528/552 (96%)	509 (96%)	17 (3%)	2 (0%)	30	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	397	ALA
1	A	276	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/241 (93%)	212 (95%)	12 (5%)	20	25
1	B	222/241 (92%)	214 (96%)	8 (4%)	31	42
All	All	446/482 (92%)	426 (96%)	20 (4%)	24	33

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	245	LYS
1	A	248	LEU
1	A	263	LYS
1	A	273	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	279	GLU
1	A	294	LYS
1	A	304	VAL
1	A	319	THR
1	A	357	LYS
1	A	358	ASN
1	A	467	LYS
1	A	494	GLU
1	B	238	GLU
1	B	263	LYS
1	B	267	THR
1	B	273	LEU
1	B	304	VAL
1	B	313	ILE
1	B	391	ASP
1	B	444	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	300	GLN
1	B	491	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PTR	A	393	1	15,16,17	1.18	1 (6%)	17,22,24	0.65	0
1	PTR	B	393	1	15,16,17	1.31	1 (6%)	17,22,24	0.76	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	393	1	-	1/10/11/13	0/1/1/1
1	PTR	B	393	1	-	0/10/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	393	PTR	OH-CZ	-4.15	1.31	1.40
1	B	393	PTR	OH-CZ	-4.08	1.31	1.40

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	393	PTR	O3P-P-OH	2.05	111.39	105.32

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	393	PTR	C-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	393	PTR	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	8IW	B	601	-	33,33,34	2.65	15 (45%)	47,48,49	1.95	16 (34%)
2	8IW	A	601	-	33,33,34	2.53	15 (45%)	47,48,49	2.09	17 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8IW	B	601	-	-	0/18/18/20	0/4/4/4
2	8IW	A	601	-	-	0/18/18/20	0/4/4/4

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	8IW	CAG-NAH	6.11	1.43	1.34
2	A	601	8IW	CAG-NAH	6.00	1.43	1.34
2	B	601	8IW	CAQ-CAP	5.39	1.48	1.39
2	A	601	8IW	CAQ-CAP	5.22	1.48	1.39
2	A	601	8IW	OBA-CAZ	4.81	1.45	1.37
2	B	601	8IW	CAS-CAT	4.78	1.42	1.37
2	B	601	8IW	OBA-CAZ	4.50	1.44	1.37
2	B	601	8IW	CAP-CAO	-4.48	1.36	1.41
2	A	601	8IW	CAP-CAO	-4.37	1.36	1.41
2	B	601	8IW	CAS-NAR	-3.97	1.30	1.37
2	B	601	8IW	CAA-CAG	3.94	1.56	1.50
2	A	601	8IW	CAA-CAG	3.69	1.56	1.50
2	A	601	8IW	CAS-NAR	-3.68	1.31	1.37
2	A	601	8IW	CAS-CAT	3.67	1.41	1.37
2	B	601	8IW	OBE-CAX	2.94	1.42	1.36
2	A	601	8IW	CAO-NAR	2.92	1.41	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	8IW	OAK-CAG	2.74	1.28	1.22
2	B	601	8IW	CAX-CAW	2.68	1.43	1.40
2	B	601	8IW	OAK-CAG	2.66	1.27	1.22
2	B	601	8IW	CAO-NAR	2.61	1.40	1.36
2	B	601	8IW	CAQ-CAL	2.51	1.44	1.39
2	B	601	8IW	CAU-CAZ	2.38	1.45	1.40
2	A	601	8IW	CAL-CAE	2.37	1.54	1.49
2	A	601	8IW	OBE-CAX	2.35	1.41	1.36
2	B	601	8IW	CBC-CAW	2.34	1.55	1.51
2	B	601	8IW	CAL-CAE	2.26	1.54	1.49
2	A	601	8IW	CAU-CAZ	2.25	1.44	1.40
2	A	601	8IW	CAX-CAW	2.20	1.43	1.40
2	A	601	8IW	CAQ-CAL	2.09	1.43	1.39
2	A	601	8IW	CAV-CAU	-2.05	1.36	1.39

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	8IW	CAP-CAO-NAN	-5.01	120.23	126.31
2	A	601	8IW	CAQ-CAP-CAO	4.40	120.89	117.64
2	B	601	8IW	CAP-CAO-NAN	-4.30	121.10	126.31
2	B	601	8IW	CAO-NAR-CAS	3.94	110.34	108.48
2	A	601	8IW	CBB-OBA-CAZ	-3.75	112.00	117.51
2	A	601	8IW	CAA-CAG-NAH	3.53	122.96	119.10
2	B	601	8IW	CAD-NAC-CAB	3.50	122.27	117.51
2	B	601	8IW	CAA-CAG-NAH	3.46	122.88	119.10
2	B	601	8IW	CAQ-CAL-CAM	3.32	120.66	117.10
2	A	601	8IW	CAV-CAW-CAX	3.27	121.08	117.64
2	A	601	8IW	CAO-NAR-CAS	3.24	110.01	108.48
2	A	601	8IW	CAE-CAF-CAA	-3.20	117.62	121.04
2	A	601	8IW	CAD-NAC-CAB	3.05	121.65	117.51
2	B	601	8IW	CAE-CAF-CAA	-3.00	117.82	121.04
2	B	601	8IW	CAQ-CAP-CAO	2.96	119.82	117.64
2	A	601	8IW	CAO-CAP-CAT	-2.96	105.20	106.22
2	A	601	8IW	CAQ-CAL-CAM	2.82	120.12	117.10
2	B	601	8IW	CAM-NAN-CAO	2.82	121.71	116.19
2	B	601	8IW	OBA-CAZ-CAY	-2.77	119.30	124.08
2	A	601	8IW	CAM-NAN-CAO	2.73	121.55	116.19
2	A	601	8IW	CAP-CAQ-CAL	-2.68	115.83	120.82
2	B	601	8IW	CAP-CAQ-CAL	-2.59	116.01	120.82
2	B	601	8IW	OBA-CAZ-CAU	2.56	120.28	116.55
2	B	601	8IW	CAV-CAW-CAX	2.39	120.16	117.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	8IW	CAU-CAT-CAP	2.38	129.93	125.97
2	A	601	8IW	NAR-CAO-NAN	2.35	130.49	126.69
2	A	601	8IW	CAP-CAO-NAR	2.34	109.28	108.42
2	A	601	8IW	OBA-CAZ-CAY	-2.31	120.10	124.08
2	B	601	8IW	CAL-CAM-NAN	-2.25	120.70	124.28
2	A	601	8IW	OBA-CAZ-CAU	2.13	119.64	116.55
2	B	601	8IW	CAA-CAB-NAC	-2.12	120.36	123.50
2	A	601	8IW	CAT-CAS-NAR	-2.02	109.16	110.46
2	B	601	8IW	CBB-OBA-CAZ	-2.01	114.57	117.51

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	8IW	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/276 (97%)	-0.21	14 (5%) 33 29	16, 29, 62, 84	0
1	B	266/276 (96%)	0.09	21 (7%) 18 16	20, 35, 82, 99	0
All	All	534/552 (96%)	-0.06	35 (6%) 24 21	16, 32, 75, 99	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	232	TYR	6.2
1	A	233	ASP	4.5
1	A	249	GLY	4.5
1	B	390	GLY	4.5
1	B	278	MET	4.2
1	A	276	ASP	4.2
1	A	232	TYR	4.0
1	A	277	THR	4.0
1	A	390	GLY	4.0
1	B	279	GLU	3.8
1	B	397	ALA	3.7
1	B	233	ASP	3.5
1	A	307	ARG	3.4
1	B	275	GLU	3.4
1	A	389	THR	3.3
1	B	307	ARG	3.3
1	B	274	LYS	3.0
1	B	401	PHE	2.8
1	B	246	HIS	2.7
1	A	500	SER	2.6
1	B	308	GLU	2.6
1	B	239	ARG	2.5
1	B	396	HIS	2.5
1	A	275	GLU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	306	THR	2.4
1	A	250	GLY	2.3
1	B	309	PRO	2.3
1	A	238	GLU	2.3
1	B	280	VAL	2.3
1	A	381	ASP	2.2
1	B	273	LEU	2.2
1	B	253	TYR	2.1
1	B	248	LEU	2.1
1	B	240	THR	2.0
1	B	389	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PTR	B	393	16/17	0.95	0.08	30,32,66,68	0
1	PTR	A	393	16/17	0.96	0.07	25,32,56,57	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	8IW	B	601	30/31	0.93	0.10	26,35,73,76	0
2	8IW	A	601	30/31	0.95	0.10	18,26,74,86	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.




Electron density around 8IW B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 8IW A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.