



Full wwPDB EM Validation Report ⓘ

Mar 8, 2026 – 05:37 PM UTC

PDB ID : 7WWK / pdb_00007wwk
EMDB ID : EMD-32868
Title : Local refinement of the SARS-CoV-2 BA.1 Spike trimer in complex with 55A8 Fab
Authors : Guo, H.; Gao, Y.; Lu, Y.; Yang, H.; Ji, X.
Deposited on : 2022-02-13
Resolution : 3.40 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14717 (2.90 - 3.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	194	11% 85% 15%
1	C	194	11% 82% 18%
2	F	108	85% 15%
2	H	108	91% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	121	 7% 74% 26%
3	I	121	 1% 82% 18%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6620 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	B	193	Total	C	N	O	S	0	0
			1550	1001	261	280	8		
1	C	194	Total	C	N	O	S	0	0
			1558	1005	263	282	8		

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	339	ASP	GLY	variant	UNP P0DTC2
B	371	LEU	SER	variant	UNP P0DTC2
B	373	PRO	SER	variant	UNP P0DTC2
B	375	PHE	SER	variant	UNP P0DTC2
B	417	ASN	LYS	variant	UNP P0DTC2
B	440	LYS	ASN	variant	UNP P0DTC2
B	446	SER	GLY	variant	UNP P0DTC2
B	477	ASN	SER	variant	UNP P0DTC2
B	478	LYS	THR	variant	UNP P0DTC2
B	484	ALA	GLU	variant	UNP P0DTC2
B	493	ARG	GLN	variant	UNP P0DTC2
B	496	SER	GLY	variant	UNP P0DTC2
B	498	ARG	GLN	variant	UNP P0DTC2
B	501	TYR	ASN	variant	UNP P0DTC2
B	505	HIS	TYR	variant	UNP P0DTC2
C	339	ASP	GLY	variant	UNP P0DTC2
C	371	LEU	SER	variant	UNP P0DTC2
C	373	PRO	SER	variant	UNP P0DTC2
C	375	PHE	SER	variant	UNP P0DTC2
C	417	ASN	LYS	variant	UNP P0DTC2
C	440	LYS	ASN	variant	UNP P0DTC2
C	446	SER	GLY	variant	UNP P0DTC2
C	477	ASN	SER	variant	UNP P0DTC2
C	478	LYS	THR	variant	UNP P0DTC2
C	484	ALA	GLU	variant	UNP P0DTC2
C	493	ARG	GLN	variant	UNP P0DTC2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	496	SER	GLY	variant	UNP P0DTC2
C	498	ARG	GLN	variant	UNP P0DTC2
C	501	TYR	ASN	variant	UNP P0DTC2
C	505	HIS	TYR	variant	UNP P0DTC2

- Molecule 2 is a protein called 55A8 light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	F	108	Total	C	N	O	S	0	0
			834	524	140	167	3		
2	H	108	Total	C	N	O	S	0	0
			834	524	140	167	3		


- Molecule 3 is a protein called 55A8 light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	121	Total	C	N	O	S	0	0
			922	580	156	182	4		
3	I	121	Total	C	N	O	S	0	0
			922	580	156	182	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike glycoprotein




- Molecule 1: Spike glycoprotein




- Molecule 2: 55A8 light chain




- Molecule 2: 55A8 light chain



• Molecule 3: 55A8 light chain



• Molecule 3: 55A8 light chain



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	181072	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.080	Depositor
Minimum map value	-0.659	Depositor
Average map value	-0.003	Depositor
Map value standard deviation	0.028	Depositor
Recommended contour level	0.4206	Depositor
Map size (Å)	419.84, 419.84, 419.84	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.82, 0.82, 0.82	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.34	0/1597	0.48	0/2174
1	C	0.36	1/1605 (0.1%)	0.49	0/2185
2	F	0.14	0/853	0.34	0/1155
2	H	0.16	0/853	0.37	0/1155
3	G	0.30	0/941	0.52	0/1275
3	I	0.26	0/941	0.47	0/1275
All	All	0.29	1/6790 (0.0%)	0.46	0/9219

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	392	PHE	C-N	5.07	1.40	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1550	0	1481	20	0
1	C	1558	0	1487	23	0
2	F	834	0	813	11	0
2	H	834	0	813	8	0
3	G	922	0	899	22	0
3	I	922	0	899	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6620	0	6392	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:35:ILE:HD11	3:I:108:SER:HB3	1.65	0.77
1:B:336:CYS:SG	1:B:337:PRO:HD2	2.25	0.76
1:B:335:LEU:HD23	1:B:362:VAL:HG13	1.70	0.73
1:C:383:SER:HB3	1:C:386:LYS:HG2	1.75	0.67
1:B:387:LEU:HD23	1:B:390:LEU:HD21	1.75	0.67
2:H:42:LYS:HE3	2:H:42:LYS:HA	1.77	0.67
1:C:346:ARG:NH1	1:C:347:PHE:O	2.28	0.65
1:C:403:ARG:HB3	1:C:406:GLU:HG3	1.79	0.65
3:I:32:TYR:CG	3:I:98:ARG:HD2	2.33	0.64
3:G:40:ALA:HB3	3:G:43:GLN:HG3	1.82	0.62
3:I:98:ARG:HH12	3:I:100:THR:HG22	1.64	0.62
3:G:94:TYR:O	3:G:114:GLY:HA2	2.00	0.62
3:I:104:ASP:HB3	3:I:107:VAL:HG12	1.83	0.60
3:G:74:LYS:HA	3:G:74:LYS:HE3	1.85	0.58
2:F:22:THR:HG22	2:F:72:THR:HG22	1.84	0.58
1:B:391:CYS:HB3	1:B:522:ALA:HB1	1.85	0.58
3:G:97:ALA:HB1	3:G:108:SER:HB2	1.85	0.58
1:B:411:ALA:HB3	1:B:414:GLN:HG3	1.85	0.56
2:F:7:SER:OG	2:F:22:THR:OG1	2.21	0.56
3:I:94:TYR:O	3:I:114:GLY:HA2	2.06	0.55
1:B:363:ALA:O	1:B:526:GLY:HA2	2.06	0.55
1:C:450:ASN:ND2	3:I:104:ASP:HA	2.21	0.55
1:C:450:ASN:HD21	3:I:104:ASP:HA	1.71	0.54
1:B:394:ASN:HB2	1:B:516:GLU:OE1	2.08	0.53
1:C:337:PRO:O	1:C:340:GLU:HG2	2.08	0.53
3:G:33:SER:HB3	3:G:105:TYR:CD1	2.44	0.52
1:C:336:CYS:HB2	1:C:338:PHE:CE2	2.44	0.52
3:G:32:TYR:CG	3:G:98:ARG:HD2	2.44	0.52
1:C:418:ILE:HA	1:C:422:ASN:HD22	1.73	0.51
3:I:12:LYS:NZ	3:I:17:SER:O	2.38	0.51
3:G:94:TYR:HE1	3:G:117:VAL:HB	1.76	0.51
1:C:362:VAL:HG22	1:C:527:PRO:HD3	1.94	0.50
3:G:51:ILE:HD13	3:G:70:ILE:HG22	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:104:ASP:HB3	3:G:107:VAL:HG22	1.92	0.50
1:B:390:LEU:HD13	1:B:392:PHE:CE2	2.48	0.49
1:C:338:PHE:O	1:C:342:PHE:HD1	1.95	0.49
1:C:393:THR:HA	1:C:522:ALA:HA	1.94	0.49
1:C:523:THR:HG23	1:C:524:VAL:HG23	1.96	0.48
1:C:526:GLY:O	1:C:527:PRO:C	2.57	0.48
2:F:2:ILE:HD13	2:F:29:ILE:HG22	1.96	0.48
3:G:26:GLY:O	3:G:32:TYR:OH	2.24	0.48
3:G:48:MET:HE1	3:G:94:TYR:CE2	2.49	0.48
1:C:446:SER:HB2	2:H:94:TYR:CE1	2.49	0.47
2:F:39:LYS:HD3	2:F:84:ALA:HB2	1.96	0.47
3:G:48:MET:HE1	3:G:94:TYR:CD2	2.49	0.47
3:G:86:LEU:HB3	3:G:119:VAL:HG11	1.97	0.47
1:C:401:VAL:HG22	1:C:509:ARG:HG2	1.96	0.47
1:C:391:CYS:HB3	1:C:522:ALA:CB	2.45	0.46
1:C:362:VAL:CG2	1:C:527:PRO:HD3	2.46	0.46
1:B:391:CYS:HB3	1:B:522:ALA:CB	2.45	0.46
1:B:516:GLU:N	1:B:516:GLU:OE2	2.49	0.46
1:C:335:LEU:HA	1:C:362:VAL:O	2.16	0.45
1:C:390:LEU:HD23	1:C:392:PHE:CE1	2.51	0.45
3:I:35:ILE:HG22	3:I:50:ARG:HG3	1.98	0.45
1:B:433:VAL:HG12	1:B:512:VAL:HG22	1.97	0.45
3:I:67:ARG:NH2	3:I:90:ASP:OD2	2.50	0.45
3:G:32:TYR:CD2	3:G:98:ARG:HD2	2.52	0.44
3:G:101:GLU:HB2	3:G:105:TYR:OH	2.17	0.44
2:F:94:TYR:OH	3:G:59:ASN:ND2	2.49	0.44
2:H:18:ARG:HG3	2:H:76:SER:HA	1.99	0.44
1:B:358:ILE:HB	1:B:395:VAL:O	2.18	0.44
3:G:39:GLN:HB2	3:G:45:LEU:HD23	1.99	0.44
1:B:393:THR:HG22	1:B:516:GLU:O	2.18	0.43
1:B:384:PRO:HA	1:B:387:LEU:HD12	2.00	0.43
2:F:94:TYR:HE1	3:G:50:ARG:HH12	1.66	0.43
2:H:107:LYS:HB2	2:H:107:LYS:HE3	1.82	0.43
1:B:355:ARG:HA	1:B:355:ARG:HD3	1.89	0.43
1:B:480:CYS:HB2	1:B:483:VAL:HG12	1.99	0.43
2:H:104:LEU:HD12	2:H:104:LEU:HA	1.91	0.43
1:B:386:LYS:HG3	1:B:389:ASP:OD2	2.19	0.43
2:H:39:LYS:HD3	2:H:84:ALA:HB2	2.01	0.43
1:B:450:ASN:ND2	3:G:104:ASP:HA	2.34	0.43
2:F:4:MET:HE3	2:F:4:MET:HB3	1.84	0.43
1:C:376:THR:O	1:C:434:ILE:HA	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:33:SER:HB2	3:I:52:ILE:HD13	2.02	0.42
2:F:46:LEU:HD23	2:F:55:GLU:HG3	2.00	0.42
2:H:2:ILE:H	2:H:2:ILE:HD12	1.85	0.42
3:I:54:ILE:HD12	3:I:54:ILE:HA	1.87	0.42
3:I:62:GLN:HA	3:I:62:GLN:OE1	2.20	0.42
2:F:83:PHE:CZ	2:F:106:ILE:HG22	2.54	0.42
3:G:6:GLN:HE21	3:G:115:THR:HG23	1.85	0.42
2:H:3:GLN:HA	2:H:3:GLN:OE1	2.20	0.42
3:I:32:TYR:CD2	3:I:98:ARG:HD2	2.54	0.42
2:F:29:ILE:HA	2:F:92:ASN:HD22	1.85	0.42
1:B:405:ASP:OD1	1:B:405:ASP:N	2.53	0.41
1:C:353:TRP:CD1	1:C:353:TRP:H	2.38	0.41
1:C:480:CYS:O	1:C:483:VAL:HG22	2.21	0.41
3:G:91:THR:O	3:G:92:ALA:HB2	2.20	0.41
3:I:35:ILE:HG13	3:I:97:ALA:HB3	2.03	0.41
3:I:89:GLU:OE1	3:I:89:GLU:N	2.36	0.41
1:C:411:ALA:HB3	1:C:414:GLN:HG3	2.02	0.41
3:I:87:ARG:HH11	3:I:87:ARG:HG2	1.86	0.40
2:F:11:LEU:HD12	2:F:12:SER:H	1.87	0.40
1:B:425:LEU:HD23	1:B:425:LEU:HA	1.85	0.40
3:G:91:THR:OG1	3:G:118:THR:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	191/194 (98%)	187 (98%)	4 (2%)	0	100	100
1	C	192/194 (99%)	184 (96%)	8 (4%)	0	100	100
2	F	106/108 (98%)	97 (92%)	9 (8%)	0	100	100
2	H	106/108 (98%)	98 (92%)	8 (8%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	G	119/121 (98%)	116 (98%)	2 (2%)	1 (1%)	16	44
3	I	119/121 (98%)	115 (97%)	3 (2%)	1 (1%)	16	44
All	All	833/846 (98%)	797 (96%)	34 (4%)	2 (0%)	44	71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	I	103	GLY
3	G	103	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	168/169 (99%)	168 (100%)	0	100	100
1	C	169/169 (100%)	169 (100%)	0	100	100
2	F	94/94 (100%)	94 (100%)	0	100	100
2	H	94/94 (100%)	94 (100%)	0	100	100
3	G	99/99 (100%)	99 (100%)	0	100	100
3	I	99/99 (100%)	99 (100%)	0	100	100
All	All	723/724 (100%)	723 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	519	HIS
2	F	37	GLN
2	F	38	GLN
3	I	109	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-32868. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 256



Y Index: 256




Z Index: 256


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 257



Y Index: 238




Z Index: 168


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views

This section was not generated.


6.6 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 50 nm³; this corresponds to an approximate mass of 45 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8 Fourier-Shell correlation ⓘ

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-32868 and PDB model 7WWK. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)

This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 81% of all backbone atoms, 72% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.4206) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7180	0.5180
B	0.6910	0.5170
C	0.6940	0.5110
F	0.7550	0.5160
G	0.6940	0.5020
H	0.7920	0.5390
I	0.7240	0.5300

