



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:03 PM UTC

PDB ID : 6X8E / pdb_00006x8e
Title : Crystal structure of JAK2 with Compound 11
Authors : Vajdos, F.F.; Knafels, J.D.
Deposited on : 2020-06-01
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	 85% 7% 8%
1	B	298	 84% 10% 6%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5246 atoms, of which 36 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase JAK2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	274	Total	C	N	O	P	S	0	1	0
			2293	1459	397	422	2	13			
1	B	281	Total	C	N	O	P	S	0	1	0
			2334	1480	402	436	2	14			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	835	GLY	-	expression tag	UNP O60674
A	836	SER	-	expression tag	UNP O60674
A	1073	SER	MET	engineered mutation	UNP O60674
A	1076	THR	PHE	engineered mutation	UNP O60674
B	835	GLY	-	expression tag	UNP O60674
B	836	SER	-	expression tag	UNP O60674
B	1073	SER	MET	engineered mutation	UNP O60674
B	1076	THR	PHE	engineered mutation	UNP O60674

- Molecule 2 is [3-{4-[6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyrazin-4-yl]-1H-pyrazol-1-yl}-1-(2,2,2-trifluoroethyl)azetidin-3-yl]acetonitrile (CCD ID: UWP) (formula: C₂₀H₁₈F₃N₉) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 50	C 20	F 3	H 18	N 9	18	0
2	B	1	Total 50	C 20	F 3	H 18	N 9	18	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	223	Total	O	0	0
			223	223		
3	B	296	Total	O	0	0
			296	296		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosine-protein kinase JAK2



• Molecule 1: Tyrosine-protein kinase JAK2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	36.72Å 153.75Å 53.07Å 90.00° 110.21° 90.00°	Depositor
Resolution (Å)	76.88 – 1.75 76.88 – 1.75	Depositor EDS
% Data completeness (in resolution range)	82.0 (76.88-1.75) 82.0 (76.88-1.75)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 1.75Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.181 , 0.225 0.194 , 0.240	Depositor DCC
R_{free} test set	2308 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtriage
Anisotropy	0.456	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.133 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5246	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UWP, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	2/2308 (0.1%)	1.19	2/3103 (0.1%)
1	B	0.85	0/2350	1.18	5/3158 (0.2%)
All	All	0.85	2/4658 (0.0%)	1.19	7/6261 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1102	MET	SD-CE	-7.09	1.61	1.79
1	A	1098	ILE	CG1-CD1	5.75	1.74	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1092	ASP	CA-CB-CG	5.74	118.34	112.60
1	A	1048	PHE	CA-CB-CG	-5.47	108.33	113.80
1	B	845	GLU	CA-C-N	5.43	127.87	120.54
1	B	845	GLU	C-N-CA	5.43	127.87	120.54
1	B	1092	ASP	CA-CB-CG	5.39	117.99	112.60
1	B	1006	GLU	CB-CG-CD	5.38	121.75	112.60
1	B	1048	PHE	CA-CB-CG	-5.17	108.63	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2293	0	2280	7	0
1	B	2334	0	2310	9	0
2	A	32	18	0	0	0
2	B	32	18	0	0	0
3	A	223	0	0	1	0
3	B	296	0	0	1	0
All	All	5210	36	4590	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:979:THR:HA	1:A:982:ILE:HD12	1.86	0.57
1:B:848:HIS:HD2	1:B:872:GLN:HE21	1.53	0.56
1:B:854:GLN:HE22	1:B:862:SER:HB2	1.73	0.54
1:B:979:THR:HA	1:B:982:ILE:HD12	1.89	0.54
1:A:1060:GLU:HG2	1:A:1089:PRO:CD	2.41	0.51
1:A:915:GLY:HA3	3:A:4168:HOH:O	2.13	0.49
1:B:917:CYS:HB3	1:B:926:LYS:HB2	1.97	0.47
1:A:980:ARG:HH11	1:A:980:ARG:HG2	1.80	0.46
1:B:1045:TYR:HB2	1:B:1102[A]:MET:HE3	1.97	0.45
1:B:1060:GLU:HG2	1:B:1089:PRO:CD	2.46	0.45
1:B:949:ASP:HB2	3:B:4231:HOH:O	2.16	0.45
1:A:1102:MET:HE3	1:A:1106:TRP:HH2	1.82	0.44
1:A:1102:MET:HE3	1:A:1106:TRP:CH2	2.54	0.43
1:B:848:HIS:CD2	1:B:872:GLN:HE21	2.33	0.43
1:B:1060:GLU:HG3	1:B:1064:MET:HE2	2.00	0.43
1:A:1060:GLU:HG2	1:A:1089:PRO:HD2	2.02	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/298 (89%)	257 (97%)	7 (3%)	1 (0%)	30	15
1	B	274/298 (92%)	269 (98%)	5 (2%)	0	100	100
All	All	539/596 (90%)	526 (98%)	12 (2%)	1 (0%)	43	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	873	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/269 (94%)	246 (97%)	7 (3%)	38	18
1	B	257/269 (96%)	246 (96%)	11 (4%)	26	8
All	All	510/538 (95%)	492 (96%)	18 (4%)	32	12

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	850	LYS
1	A	947	ARG
1	A	977	LEU
1	A	985	GLU
1	A	1010	VAL
1	A	1083	LYS
1	A	1121	LEU
1	B	871	LEU
1	B	887	SER
1	B	937	LEU
1	B	947	ARG
1	B	977	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	985	GLU
1	B	1006	GLU
1	B	1010	VAL
1	B	1083	LYS
1	B	1121	LEU
1	B	1130	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	854	GLN
1	A	885	GLN
1	A	886	HIS
1	A	909	ASN
1	A	924	ASN
1	A	950	HIS
1	A	988	ASN
1	A	1111	ASN
1	A	1112	GLN
1	B	848	HIS
1	B	854	GLN
1	B	885	GLN
1	B	909	ASN
1	B	955	GLN
1	B	988	ASN
1	B	1070	GLN
1	B	1072	GLN
1	B	1085	ASN
1	B	1111	ASN
1	B	1112	GLN
1	B	1125	GLN
1	B	1129	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	PTR	A	1008	1	15,16,17	1.93	5 (33%)	17,22,24	1.09	1 (5%)
1	PTR	A	1007	1	15,16,17	1.92	5 (33%)	17,22,24	1.21	2 (11%)
1	PTR	B	1008	1	15,16,17	1.78	4 (26%)	17,22,24	1.97	6 (35%)
1	PTR	B	1007	1	15,16,17	1.75	3 (20%)	17,22,24	1.38	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	1008	1	-	0/10/11/13	0/1/1/1
1	PTR	A	1007	1	-	1/10/11/13	0/1/1/1
1	PTR	B	1008	1	-	0/10/11/13	0/1/1/1
1	PTR	B	1007	1	-	0/10/11/13	0/1/1/1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1008	PTR	CE1-CZ	4.90	1.47	1.38
1	B	1007	PTR	CE2-CZ	3.74	1.45	1.38
1	A	1007	PTR	CE1-CZ	3.40	1.45	1.38
1	B	1007	PTR	P-OH	-3.35	1.53	1.59
1	A	1007	PTR	CE1-CD1	3.33	1.44	1.38
1	B	1008	PTR	CD1-CG	3.23	1.45	1.38
1	A	1007	PTR	CD1-CG	3.20	1.45	1.38
1	B	1007	PTR	CD2-CG	2.98	1.44	1.38
1	B	1008	PTR	CE1-CZ	2.75	1.43	1.38
1	B	1008	PTR	CB-CA	2.61	1.59	1.53
1	A	1008	PTR	P-OH	-2.61	1.54	1.59
1	B	1008	PTR	CE2-CD2	2.45	1.42	1.38
1	A	1008	PTR	CD2-CG	2.31	1.43	1.38
1	A	1007	PTR	CD2-CG	2.30	1.43	1.38
1	A	1007	PTR	P-OH	-2.27	1.55	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1008	PTR	CE1-CD1	2.09	1.42	1.38
1	A	1008	PTR	CE2-CD2	2.02	1.42	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1008	PTR	O3P-P-OH	3.91	116.86	105.32
1	B	1008	PTR	CB-CG-CD2	-3.07	115.19	120.90
1	B	1008	PTR	OH-CZ-CE1	2.80	127.62	119.22
1	B	1008	PTR	OH-CZ-CE2	-2.64	111.31	119.22
1	B	1008	PTR	O3P-P-O2P	2.59	117.51	107.80
1	A	1007	PTR	O2P-P-OH	2.58	112.96	105.32
1	B	1007	PTR	CB-CG-CD1	-2.53	116.19	120.90
1	A	1007	PTR	CE1-CD1-CG	-2.30	117.97	121.00
1	B	1008	PTR	O2P-P-OH	-2.26	98.65	105.32
1	B	1007	PTR	O2P-P-OH	2.07	111.43	105.32
1	A	1008	PTR	OH-CZ-CE1	2.03	125.29	119.22

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1007	PTR	CZ-OH-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	UWP	B	4001	-	34,36,36	1.84	6 (17%)	34,55,55	1.42	8 (23%)
2	UWP	A	4000	-	34,36,36	1.80	8 (23%)	34,55,55	1.49	9 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UWP	B	4001	-	-	0/14/33/33	0/5/5/5
2	UWP	A	4000	-	-	0/14/33/33	0/5/5/5

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	4001	UWP	C2-N1	4.72	1.39	1.34
2	B	4001	UWP	C13-N7	4.67	1.39	1.35
2	A	4000	UWP	C2-N1	4.60	1.39	1.34
2	B	4001	UWP	C17-C18	4.59	1.58	1.50
2	A	4000	UWP	C17-C18	4.51	1.58	1.50
2	A	4000	UWP	C13-N7	4.19	1.39	1.35
2	B	4001	UWP	C15-N8	2.52	1.51	1.49
2	B	4001	UWP	C16-N8	2.27	1.50	1.49
2	A	4000	UWP	C16-N8	2.18	1.50	1.49
2	A	4000	UWP	C4-N2	2.17	1.36	1.32
2	A	4000	UWP	C15-N8	2.17	1.50	1.49
2	B	4001	UWP	C4-N2	2.12	1.36	1.32
2	A	4000	UWP	C15-C14	2.11	1.57	1.55
2	A	4000	UWP	C6-N3	2.07	1.39	1.36

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	4000	UWP	C18-C17-N8	-3.75	104.50	112.43
2	B	4001	UWP	C18-C17-N8	-3.53	104.95	112.43
2	A	4000	UWP	C13-C11-C12	2.50	106.43	104.22
2	A	4000	UWP	C11-C13-N7	-2.44	105.50	107.75
2	B	4001	UWP	C13-C11-C12	2.27	106.22	104.22
2	B	4001	UWP	C3-C4-N2	-2.26	110.30	112.09

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4001	UWP	C11-C13-N7	-2.24	105.68	107.75
2	A	4000	UWP	C1-N1-C2	-2.23	125.70	127.94
2	A	4000	UWP	C11-C12-N6	-2.19	110.35	112.09
2	A	4000	UWP	C1-N1-N2	2.18	121.78	120.03
2	B	4001	UWP	C2-C3-C4	2.18	106.14	104.22
2	A	4000	UWP	C3-C4-N2	-2.15	110.38	112.09
2	B	4001	UWP	C3-C2-N1	-2.10	105.61	107.22
2	A	4000	UWP	C2-C3-C4	2.09	106.07	104.22
2	B	4001	UWP	C11-C12-N6	-2.06	110.46	112.09
2	A	4000	UWP	C3-C2-N1	-2.03	105.66	107.22
2	B	4001	UWP	C4-N2-N1	2.02	105.52	104.41


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/298 (91%)	-1.11	0 100 100	9, 24, 63, 82	1 (0%)
1	B	279/298 (93%)	-1.21	0 100 100	8, 20, 52, 79	1 (0%)
All	All	551/596 (92%)	-1.16	0 100 100	8, 22, 58, 82	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PTR	A	1007	16/17	0.99	0.03	17,26,42,43	0
1	PTR	A	1008	16/17	0.99	0.04	19,28,41,41	0
1	PTR	B	1007	16/17	0.99	0.04	22,32,47,48	0
1	PTR	B	1008	16/17	1.00	0.02	14,19,26,26	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	UWP	A	4000	32/32	0.99	0.03	13,21,35,41	18
2	UWP	B	4001	32/32	0.99	0.03	13,22,35,40	18

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around UWP A 4000:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UWP B 4001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.