wwPDB EM Validation Summary Report (i)

Oct 19, 2026 – 09:38 PM JST

PDB ID	:	9X8O / pdb_00009x8o
EMDB ID	:	EMD-66656
Title	:	Structure of Candida aurisglycosylphosphatidylinositol transamidase
Authors	:	Ding, X.Y.; Hua, Z.K.; Zhang, M.J.; Zhang, M.; Liu, X.T.; Yu, H.J.
Deposited on	:	2025-10-19
Resolution	:	3.19 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev133
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4-5-2 with Phenix2.2
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14455 (2.69 - 3.69)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	812	54% 11% 34%
2	C	834	60% 9% 31%
3	D	658	53% 12% 35%
4	E	733	49% 15% 36%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	F	659	

2 Entry composition (i)

There are 14 unique types of molecules in this entry. The entry contains 18688 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gpi16 subunit,Fusion protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	532	Total	C	N	O	S	0	0
			4208	2708	715	770	15		

- Molecule 2 is a protein called Gaa1,Fusion protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	577	Total	C	N	O	S	0	0
			4496	2928	741	809	18		

- Molecule 3 is a protein called Gab1,Fusion protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	426	Total	C	N	O	S	0	0
			3432	2308	525	584	15		


- Molecule 4 is a protein called GPI transamidase component PIG-S,Fusion protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	468	Total	C	N	O	S	0	0
			3687	2373	603	693	18		

- Molecule 5 is a protein called GPI-anchor transamidase,Fusion protein.


Mol	Chain	Residues	Atoms					AltConf	Trace
5	F	312	Total	C	N	O	S	0	0
			2516	1601	426	478	11		

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms	AltConf
6	A	1	Total C N O 14 8 1 5	0
6	A	1	Total C N O 14 8 1 5	0
6	A	1	Total C N O 14 8 1 5	0
6	F	1	Total C N O 14 8 1 5	0

- Molecule 7 is [(2 {R})-3-[[(1 {R},2 {S},3 {R},4 {R},5 {S},6 {R})-2-[(2 {R},3 {R},4 {R},5 {R},6 {R})-3-azanyl-5-[(2 {R},3 {S},4 {S},5 {S},6 {R})-3-[2-azanylethoxy(oxidanyl)phosphoryl]oxy-6-(hydroxymethyl)-4,5-bis(oxidanyl)oxan-2-yl]oxy-6-(hydroxymethyl)-4-oxidanyl-oxan-2-yl]oxy-6-heptanoyloxy-3,4,5-tris(oxidanyl)cyclohexyl]oxy-oxidanyl-phosphoryl]oxy-2-butanoyloxy-propyl] heptanoate (CCD ID: A1EZD) (formula: $C_{41}H_{76}N_2O_{26}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
7	A	1	Total	C	N	O	P	0
			71	41	2	26	2	

- Molecule 8 is HEPTANE (CCD ID: HP6) (formula: C_7H_{16}).




Mol	Chain	Residues	Atoms		AltConf
8	C	1	Total	C	0
			7	7	
8	C	1	Total	C	0
			7	7	
8	C	1	Total	C	0
			7	7	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	AltConf
8	D	1	Total C 7 7	0
8	D	1	Total C 7 7	0

- Molecule 9 is DODECANE (CCD ID: D12) (formula: $C_{12}H_{26}$).




Mol	Chain	Residues	Atoms	AltConf
9	C	1	Total C 12 12	0
9	D	1	Total C 12 12	0

- Molecule 10 is TETRADECANE (CCD ID: C14) (formula: $C_{14}H_{30}$).



Mol	Chain	Residues	Atoms	AltConf
10	C	1	Total C 14 14	0

- Molecule 11 is (11R,14S)-17-amino-14-hydroxy-8,14-dioxo-9,13,15-trioxa-14lambda 5 -phosphaheptadecan-11-yl decanoate (CCD ID: XKP) (formula: $C_{23}H_{46}NO_8P$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	AltConf
11	C	1	Total C N O P 27 17 1 8 1	0
11	C	1	Total C N O P 33 23 1 8 1	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					AltConf
11	D	1	Total	C	N	O	P	0
			33	23	1	8	1	

- Molecule 12 is nonane (CCD ID: DD9) (formula: C_9H_{20}).



Mol	Chain	Residues	Atoms		AltConf
12	D	1	Total	C	0
			9	9	
12	D	1	Total	C	0
			9	9	
12	D	1	Total	C	0
			9	9	

- Molecule 13 is CHOLESTEROL (CCD ID: CLR) (formula: $C_{27}H_{46}O$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
13	D	1	Total	C	O	0
			28	27	1	

- Molecule 14 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		AltConf
14	F	1	Total	Ca	0
			1	1	

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gpi16 subunit,Fusion protein


Chain A:



- Molecule 2: Gaa1,Fusion protein

Chain C:






• Molecule 3: Gab1,Fusion protein

Chain D:







• Molecule 4: GPI transamidase component PIG-S,Fusion protein

Chain E:








• Molecule 5: GPI-anchor transamidase,Fusion protein

Chain F:





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	170112	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS TALOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50	Depositor
Minimum defocus (nm)	1100	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	3.569	Depositor
Minimum map value	-2.080	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.073	Depositor
Recommended contour level	0.169	Depositor
Map size (Å)	342.40002, 342.40002, 342.40002	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.07, 1.07, 1.07	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XKP, A1EZD, NAG, HP6, CLR, C14, DD9, CA, D12

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.13	0/4328	0.29	0/5901
2	C	0.12	0/4605	0.26	0/6275
3	D	0.13	0/3536	0.28	0/4830
4	E	0.14	0/3774	0.29	0/5134
5	F	0.12	0/2575	0.29	0/3498
All	All	0.13	0/18818	0.28	0/25638

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4208	0	4146	60	0
2	C	4496	0	4554	49	0
3	D	3432	0	3468	54	0
4	E	3687	0	3579	65	0
5	F	2516	0	2388	49	0
6	A	42	0	39	0	0
6	F	14	0	13	0	0
7	A	71	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	C	21	0	48	0	0
8	D	14	0	32	0	0
9	C	12	0	26	0	0
9	D	12	0	26	1	0
10	C	14	0	30	0	0
11	C	60	0	0	0	0
11	D	33	0	0	1	0
12	D	27	0	60	0	0
13	D	28	0	46	1	0
14	F	1	0	0	0	0
All	All	18688	0	18455	255	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 255 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:319:GLU:HG3	4:E:321:PRO:HD2	1.67	0.77
4:E:36:THR:HG22	4:E:37:THR:H	1.58	0.68
4:E:201:ARG:NH2	4:E:324:ASP:O	2.28	0.67
1:A:544:LYS:HE2	3:D:425:GLN:HB3	1.75	0.66
1:A:201:ARG:NH1	4:E:235:TYR:O	2.28	0.66

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	530/812 (65%)	497 (94%)	32 (6%)	1 (0%)	43 73
2	C	575/834 (69%)	553 (96%)	22 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	D	424/658 (64%)	415 (98%)	9 (2%)	0	100	100
4	E	464/733 (63%)	431 (93%)	33 (7%)	0	100	100
5	F	308/659 (47%)	291 (94%)	17 (6%)	0	100	100
All	All	2301/3696 (62%)	2187 (95%)	113 (5%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	167	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	462/697 (66%)	462 (100%)	0	100	100
2	C	487/700 (70%)	487 (100%)	0	100	100
3	D	377/570 (66%)	377 (100%)	0	100	100
4	E	406/653 (62%)	405 (100%)	1 (0%)	87	89
5	F	273/574 (48%)	273 (100%)	0	100	100
All	All	2005/3194 (63%)	2004 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	E	121	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
4	E	194	ASN
4	E	280	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	E	232	GLN
4	E	345	GLN
2	C	501	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 21 ligands modelled in this entry, 1 is monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
8	HP6	D	706	-	6,6,6	0.30	0	5,5,5	0.32	0
10	C14	C	904	-	13,13,13	0.28	0	12,12,12	0.38	0
6	NAG	A	903	1	14,14,15	3.04	8 (57%)	17,19,21	1.22	2 (11%)
6	NAG	F	701	5	14,14,15	3.12	8 (57%)	17,19,21	1.39	3 (17%)
13	CLR	D	707	-	31,31,31	0.37	0	48,48,48	1.03	2 (4%)
8	HP6	C	901	-	6,6,6	0.31	0	5,5,5	0.26	0
12	DD9	D	701	-	8,8,8	0.29	0	7,7,7	0.34	0
6	NAG	A	901	1	14,14,15	3.11	10 (71%)	17,19,21	1.39	3 (17%)
6	NAG	A	902	1	14,14,15	3.01	7 (50%)	17,19,21	1.39	4 (23%)
8	HP6	D	704	-	6,6,6	0.30	0	5,5,5	0.28	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
7	A1EZD	A	904	-	73,73,73	0.39	0	97,101,101	0.83	4 (4%)
11	XKP	C	906	-	26,26,32	1.18	4 (15%)	29,31,37	1.07	2 (6%)
9	D12	D	703	-	11,11,11	0.29	0	10,10,10	0.36	0
8	HP6	C	905	-	6,6,6	0.30	0	5,5,5	0.27	0
11	XKP	D	708	-	32,32,32	1.06	4 (12%)	35,37,37	1.03	2 (5%)
9	D12	C	902	-	11,11,11	0.29	0	10,10,10	0.35	0
8	HP6	C	903	-	6,6,6	0.30	0	5,5,5	0.29	0
12	DD9	D	705	-	8,8,8	0.29	0	7,7,7	0.34	0
12	DD9	D	702	-	8,8,8	0.30	0	7,7,7	0.32	0
11	XKP	C	907	-	32,32,32	1.06	4 (12%)	35,37,37	1.02	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	HP6	D	706	-	-	2/4/4/4	-
10	C14	C	904	-	-	7/11/11/11	-
6	NAG	A	903	1	-	2/6/23/26	0/1/1/1
6	NAG	F	701	5	-	0/6/23/26	0/1/1/1
13	CLR	D	707	-	-	7/10/68/68	0/4/4/4
8	HP6	C	901	-	-	3/4/4/4	-
12	DD9	D	701	-	-	3/6/6/6	-
6	NAG	A	901	1	-	2/6/23/26	0/1/1/1
6	NAG	A	902	1	-	1/6/23/26	0/1/1/1
8	HP6	D	704	-	-	2/4/4/4	-
7	A1EZD	A	904	-	-	15/61/125/125	0/3/3/3
11	XKP	C	906	-	-	16/30/30/36	-
9	D12	D	703	-	-	5/9/9/9	-
8	HP6	C	905	-	-	2/4/4/4	-
11	XKP	D	708	-	-	18/36/36/36	-
9	D12	C	902	-	-	6/9/9/9	-
8	HP6	C	903	-	-	2/4/4/4	-
12	DD9	D	705	-	-	4/6/6/6	-
12	DD9	D	702	-	-	4/6/6/6	-
11	XKP	C	907	-	-	23/36/36/36	-

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	901	NAG	O5-C5	6.20	1.56	1.43
6	F	701	NAG	O5-C5	6.15	1.55	1.43
6	A	903	NAG	O5-C5	5.87	1.55	1.43
6	A	902	NAG	O5-C5	5.68	1.55	1.43
6	A	902	NAG	C2-N2	5.68	1.56	1.46

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
11	D	708	XKP	O6-C11-C12	4.15	120.45	111.50
11	C	907	XKP	O6-C11-C12	3.99	120.09	111.50
11	C	906	XKP	O6-C11-C12	3.80	119.68	111.50
7	A	904	A1EZD	C1-C2-C3	-3.06	102.22	110.21
6	A	902	NAG	C4-C3-C2	3.00	115.42	111.02

There are no chirality outliers.

5 of 124 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	904	A1EZD	C11-C23-O22-P1
7	A	904	A1EZD	C28-C22-O20-C19
7	A	904	A1EZD	O23-C22-O20-C19
11	C	906	XKP	O1-C6-C7-O6
11	C	906	XKP	C8-O2-P-O3

There are no ring outliers.


4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
13	D	707	CLR	1	0
7	A	904	A1EZD	1	0
9	D	703	D12	1	0
11	D	708	XKP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand CLR D 707




Ligand A1EZD A 904




Ligand XKP C 906




Bond lengths



Bond angles




Torsions




Rings


Ligand XKP D 708




Bond lengths



Bond angles




Torsions



Rings

Ligand XKP C 907



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.


6 Map visualisation

This section contains visualisations of the EMDB entry EMD-66656. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



X Index: 172




Y Index: 173




Z Index: 152


6.3.2 Raw map



X Index: 154



Y Index: 173




Z Index: 152


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)


6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.169. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 404 nm^3; this corresponds to an approximate mass of 365 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.313 $Å^{-1}$

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.313 Å^{-1}

8.2 Resolution estimates (i)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.19	-	-
Author-provided FSC curve	3.19	3.70	3.23
Unmasked-calculated*	4.18	7.18	4.25

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.18 differs from the reported value 3.19 by more than 10 %


9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-66656 and PDB model 9X8O. Per-residue inclusion information can be found in section 3 on page 11.


9.1 Map-model overlay



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.169 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.


9.2 Q-score mapped to coordinate model



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.169).

9.4 Atom inclusion (i)



At the recommended contour level, 95% of all backbone atoms, 93% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.169) and Q-score for the entire model and for each chain.



Chain	Atom inclusion	Q-score
All	0.9300	0.4590
A	0.9350	0.4700
C	0.9390	0.4830
D	0.9300	0.4700
E	0.9210	0.4080
F	0.9210	0.4580