



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2026 – 07:51 AM JST

PDB ID : 5XJA / pdb_00005xja
Title : The Crystal Structure of the Minimal Core Domain of the Microtubule Depolymerizer KIF2C Complexed with ADP-Mg-AlF_x
Authors : Ogawa, T.; Jiang, X.; Hirokawa, N.
Deposited on : 2017-04-30
Resolution : 3.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1736 (3.50-3.38)
Clashscore	190562	1808 (3.50-3.38)
Ramachandran outliers	187476	1776 (3.50-3.38)
Sidechain outliers	187428	1777 (3.50-3.38)
RSRZ outliers	180081	1736 (3.50-3.38)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	426	3% 70% 8% 21%
1	B	426	3% 66% 10% 22%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kinesin-like protein KIF2C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	337	Total	C	N	O	S	0	0	0
			2650	1673	465	491	21			
1	B	332	Total	C	N	O	S	0	0	0
			2617	1653	459	484	21			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	167	MET	-	initiating methionine	UNP Q922S8
A	168	ALA	-	expression tag	UNP Q922S8
A	169	SER	-	expression tag	UNP Q922S8
A	170	MET	-	expression tag	UNP Q922S8
A	171	THR	-	expression tag	UNP Q922S8
A	172	GLY	-	expression tag	UNP Q922S8
A	173	GLY	-	expression tag	UNP Q922S8
A	174	GLN	-	expression tag	UNP Q922S8
A	175	GLN	-	expression tag	UNP Q922S8
A	176	MET	-	expression tag	UNP Q922S8
A	177	GLY	-	expression tag	UNP Q922S8
A	178	ARG	-	expression tag	UNP Q922S8
A	179	ASP	-	expression tag	UNP Q922S8
A	180	PRO	-	expression tag	UNP Q922S8
A	181	ASN	-	expression tag	UNP Q922S8
A	182	SER	-	expression tag	UNP Q922S8
A	586	HIS	-	expression tag	UNP Q922S8
A	587	HIS	-	expression tag	UNP Q922S8
A	588	HIS	-	expression tag	UNP Q922S8
A	589	HIS	-	expression tag	UNP Q922S8
A	590	HIS	-	expression tag	UNP Q922S8
A	591	HIS	-	expression tag	UNP Q922S8
A	592	HIS	-	expression tag	UNP Q922S8
B	167	MET	-	initiating methionine	UNP Q922S8
B	168	ALA	-	expression tag	UNP Q922S8

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	169	SER	-	expression tag	UNP Q922S8
B	170	MET	-	expression tag	UNP Q922S8
B	171	THR	-	expression tag	UNP Q922S8
B	172	GLY	-	expression tag	UNP Q922S8
B	173	GLY	-	expression tag	UNP Q922S8
B	174	GLN	-	expression tag	UNP Q922S8
B	175	GLN	-	expression tag	UNP Q922S8
B	176	MET	-	expression tag	UNP Q922S8
B	177	GLY	-	expression tag	UNP Q922S8
B	178	ARG	-	expression tag	UNP Q922S8
B	179	ASP	-	expression tag	UNP Q922S8
B	180	PRO	-	expression tag	UNP Q922S8
B	181	ASN	-	expression tag	UNP Q922S8
B	182	SER	-	expression tag	UNP Q922S8
B	586	HIS	-	expression tag	UNP Q922S8
B	587	HIS	-	expression tag	UNP Q922S8
B	588	HIS	-	expression tag	UNP Q922S8
B	589	HIS	-	expression tag	UNP Q922S8
B	590	HIS	-	expression tag	UNP Q922S8
B	591	HIS	-	expression tag	UNP Q922S8
B	592	HIS	-	expression tag	UNP Q922S8

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
3	B	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 4 is ALUMINUM FLUORIDE (CCD ID: AF3) (formula: AlF₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Al	F	0	0
			4	1	3		
4	B	1	Total	Al	F	0	0
			4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	4	Total 4	O 4	0	0
5	B	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Kinesin-like protein KIF2C



• Molecule 1: Kinesin-like protein KIF2C



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	91.10Å 166.99Å 74.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.43 30.00 – 3.43	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-3.43) 99.8 (30.00-3.43)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 3.47Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.198 , 0.230 0.201 , 0.237	Depositor DCC
R_{free} test set	846 reflections (5.30%)	wwPDB-VP
Wilson B-factor (Å ²)	64.9	Xtriage
Anisotropy	0.593	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5338	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AF3, ADP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	5/2688 (0.2%)	1.34	11/3611 (0.3%)
1	B	1.00	0/2655	1.18	9/3566 (0.3%)
All	All	1.01	5/5343 (0.1%)	1.26	20/7177 (0.3%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	504	GLN	C-O	11.20	1.36	1.23
1	A	465	ARG	CZ-NH1	5.67	1.40	1.32
1	A	465	ARG	CZ-NH2	5.33	1.40	1.33
1	A	465	ARG	NE-CZ	5.24	1.38	1.33
1	A	494	ARG	CA-C	-5.21	1.47	1.53

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	505	THR	CB-CA-C	-32.68	77.48	109.83
1	A	505	THR	N-CA-C	13.70	134.34	113.51
1	B	252	GLU	N-CA-C	10.67	133.53	110.80
1	A	504	GLN	CB-CA-C	-10.55	93.90	110.14
1	A	495	GLY	N-CA-C	-8.45	93.14	113.18
1	A	505	THR	CA-C-N	8.25	137.53	122.27
1	A	505	THR	C-N-CA	8.25	137.53	122.27
1	B	241	CYS	N-CA-C	7.91	120.80	110.43
1	A	504	GLN	N-CA-C	7.76	121.14	108.26
1	A	241	CYS	N-CA-C	7.03	125.77	110.80
1	A	487	VAL	N-CA-C	6.57	117.36	108.17
1	B	464	SER	CB-CA-C	-6.37	99.61	112.99
1	B	411	LYS	N-CA-C	5.96	118.47	111.02
1	B	284	LEU	N-CA-C	5.90	119.10	109.06

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	366	SER	N-CA-C	5.88	119.71	112.54
1	B	244	LEU	N-CA-C	5.72	119.73	112.87
1	B	527	GLY	N-CA-C	-5.70	99.67	113.18
1	A	550	GLY	N-CA-C	5.36	118.23	112.33
1	B	251	GLU	CB-CA-C	-5.20	100.07	110.42
1	B	384	PRO	CB-CA-C	-5.07	103.81	112.64

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2650	0	2673	42	1
1	B	2617	0	2638	34	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	27	0	12	2	0
3	B	27	0	12	0	0
4	A	4	0	0	0	0
4	B	4	0	0	0	0
5	A	4	0	0	0	0
5	B	3	0	0	0	0
All	All	5338	0	5335	70	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:505:THR:O	1:A:505:THR:HG22	1.33	1.10
1:A:504:GLN:O	1:A:505:THR:OG1	1.74	1.05
1:A:505:THR:O	1:A:505:THR:CG2	2.09	0.97
1:A:249:PRO:CB	1:A:250:ILE:HD12	2.04	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:357:ASP:HB2	1:A:367:LYS:HD3	1.57	0.86
1:A:494:ARG:C	1:A:496:ALA:H	1.86	0.83
1:A:250:ILE:HG23	1:B:380:LEU:CD1	2.09	0.82
1:A:249:PRO:C	1:A:250:ILE:HD12	2.04	0.81
1:A:249:PRO:HG2	1:A:250:ILE:HD12	1.63	0.81
1:A:249:PRO:CG	1:A:250:ILE:HD12	2.13	0.79
1:A:249:PRO:HB2	1:A:250:ILE:HD12	1.63	0.79
1:A:249:PRO:HG2	1:A:250:ILE:CD1	2.14	0.78
1:B:251:GLU:O	1:B:251:GLU:HG3	1.85	0.76
1:A:248:ASP:OD1	1:A:249:PRO:HD2	1.91	0.70
1:A:250:ILE:HG22	1:A:250:ILE:O	1.92	0.70
1:A:494:ARG:C	1:A:496:ALA:N	2.50	0.68
1:A:249:PRO:HB2	1:A:250:ILE:CD1	2.24	0.68
1:B:364:ASN:HD22	1:B:367:LYS:HB2	1.59	0.68
1:B:364:ASN:ND2	1:B:367:LYS:HB2	2.12	0.64
1:B:357:ASP:O	1:B:363:GLN:N	2.32	0.62
1:B:452:ARG:O	1:B:453:THR:OG1	2.09	0.62
1:A:350:LYS:N	3:A:602:ADP:O2B	2.32	0.62
1:A:493:GLU:O	1:A:493:GLU:HG2	2.00	0.61
1:A:367:LYS:HG2	1:A:367:LYS:O	2.00	0.61
1:A:250:ILE:HG23	1:B:380:LEU:HD13	1.84	0.60
1:A:384:PRO:HG2	1:B:254:ARG:NH1	2.17	0.59
1:B:254:ARG:NH2	1:B:551:GLU:O	2.35	0.57
1:B:420:ASP:OD1	1:B:422:ARG:N	2.38	0.56
1:B:527:GLY:O	1:B:528:GLN:CB	2.52	0.56
1:B:527:GLY:O	1:B:584:LEU:HD21	2.07	0.55
1:B:357:ASP:HB3	1:B:364:ASN:HB3	1.88	0.55
1:A:504:GLN:O	1:A:505:THR:CB	2.55	0.54
1:A:495:GLY:O	1:A:496:ALA:HB3	2.09	0.52
1:A:504:GLN:C	1:A:505:THR:OG1	2.51	0.51
1:A:452:ARG:O	1:A:452:ARG:HG2	2.10	0.51
1:A:330:GLN:OE1	1:B:249:PRO:HB2	2.11	0.50
1:A:249:PRO:O	1:A:250:ILE:HD12	2.10	0.49
1:B:326:ARG:HB3	1:B:327:PRO:HD3	1.94	0.49
1:B:401:TYR:CD2	1:B:401:TYR:C	2.90	0.49
1:B:465:ARG:HD3	1:B:508:GLU:OE2	2.13	0.49
1:A:250:ILE:HA	1:B:380:LEU:HD11	1.95	0.48
1:A:347:GLY:N	3:A:602:ADP:O3B	2.47	0.48
1:B:248:ASP:CG	1:B:248:ASP:O	2.54	0.48
1:B:411:LYS:HA	1:B:451:CYS:SG	2.55	0.47
1:A:343:TYR:CZ	1:A:577:TYR:CD2	3.04	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:442:ILE:O	1:A:446:ASN:ND2	2.49	0.45
1:A:250:ILE:HG23	1:B:380:LEU:HD11	1.92	0.45
1:B:373:ALA:O	1:B:377:VAL:HG23	2.17	0.45
1:B:557:MET:HE2	1:B:557:MET:HB2	1.82	0.44
1:A:409:LEU:HD22	1:A:444:MET:HE3	1.99	0.44
1:A:364:ASN:ND2	1:A:367:LYS:HB2	2.32	0.44
1:B:356:GLY:O	1:B:357:ASP:C	2.61	0.44
1:B:345:GLN:HG2	1:B:346:THR:O	2.18	0.44
1:A:373:ALA:O	1:A:377:VAL:HG23	2.17	0.43
1:A:249:PRO:O	1:A:250:ILE:HB	2.18	0.42
1:A:476:THR:HG23	1:A:477:LYS:N	2.34	0.42
1:B:378:PHE:CE2	1:B:437:CYS:C	2.98	0.42
1:B:523:ILE:O	1:B:526:LEU:HB3	2.20	0.42
1:B:343:TYR:CZ	1:B:577:TYR:CD2	3.07	0.42
1:B:526:LEU:HB3	1:B:527:GLY:H	1.75	0.42
1:A:249:PRO:C	1:A:250:ILE:CD1	2.84	0.42
1:A:451:CYS:SG	1:A:451:CYS:O	2.78	0.41
1:B:240:GLU:HG3	1:B:241:CYS:N	2.35	0.41
1:B:524:ARG:O	1:B:584:LEU:HD23	2.20	0.41
1:A:557:MET:HE3	1:A:581:VAL:HG21	2.02	0.41
1:B:357:ASP:HB3	1:B:364:ASN:CB	2.51	0.41
1:B:251:GLU:O	1:B:251:GLU:CG	2.64	0.41
1:A:249:PRO:CG	1:A:250:ILE:CD1	2.85	0.41
1:A:385:ARG:HD2	1:A:386:TYR:CE1	2.55	0.41
1:B:476:THR:HG23	1:B:477:LYS:N	2.36	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:GLU:OE2	1:A:301:GLU:OE2[2_655]	1.47	0.73
1:B:301:GLU:OE2	1:B:301:GLU:OE2[2_655]	1.95	0.25

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/426 (77%)	294 (90%)	32 (10%)	1 (0%)	36	66
1	B	322/426 (76%)	293 (91%)	27 (8%)	2 (1%)	21	52
All	All	649/852 (76%)	587 (90%)	59 (9%)	3 (0%)	24	56

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	250	ILE
1	B	526	LEU
1	A	250	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/374 (78%)	288 (99%)	2 (1%)	76	78
1	B	287/374 (77%)	283 (99%)	4 (1%)	59	71
All	All	577/748 (77%)	571 (99%)	6 (1%)	68	75

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	318	GLU
1	A	395	VAL
1	B	282	LYS
1	B	301	GLU
1	B	395	VAL
1	B	526	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	225	ASN
1	A	253	HIS
1	A	364	ASN
1	A	390	ASN
1	A	471	GLN
1	A	552	ASN
1	B	364	ASN
1	B	390	ASN
1	B	552	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	AF3	A	603	1,3	0,3,3	-	-	-		
4	AF3	B	603	1,3	0,3,3	-	-	-		
3	ADP	B	602	4,2	27,29,29	1.44	5 (18%)	42,45,45	2.09	12 (28%)
3	ADP	A	602	4,2	27,29,29	1.48	4 (14%)	42,45,45	1.71	6 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	B	602	4,2	-	0/16/32/32	0/3/3/3
3	ADP	A	602	4,2	-	0/16/32/32	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	ADP	C5-C4	4.45	1.47	1.39
3	A	602	ADP	C5-C4	4.42	1.47	1.39
3	B	602	ADP	C8-N7	2.82	1.36	1.31
3	A	602	ADP	C8-N7	2.80	1.36	1.31
3	A	602	ADP	C5-N7	-2.68	1.33	1.39
3	A	602	ADP	C4-N9	-2.63	1.31	1.37
3	B	602	ADP	C4-N9	-2.51	1.32	1.37
3	B	602	ADP	C5-N7	-2.22	1.34	1.39
3	B	602	ADP	C5-C6	2.20	1.47	1.41

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	ADP	C5-C4-N3	-5.49	119.59	126.75
3	B	602	ADP	PA-O3A-PB	-4.91	115.97	132.83
3	B	602	ADP	C5-C4-N3	-4.85	120.42	126.75
3	A	602	ADP	N3-C4-N9	4.44	134.40	127.08
3	B	602	ADP	O3B-PB-O2B	3.90	122.53	107.64
3	B	602	ADP	N3-C2-N1	-3.80	122.66	128.60
3	B	602	ADP	C2-N3-C4	3.73	120.56	111.75
3	B	602	ADP	N3-C4-N9	3.59	132.99	127.08
3	A	602	ADP	C2-N3-C4	3.47	119.95	111.75
3	B	602	ADP	C5'-C4'-C3'	-2.98	104.02	115.18
3	B	602	ADP	O2A-PA-O1A	2.94	126.77	112.24
3	B	602	ADP	O2'-C2'-C1'	2.89	119.68	110.02
3	A	602	ADP	N3-C2-N1	-2.82	124.20	128.60
3	B	602	ADP	O5'-PA-O1A	-2.80	98.11	109.07
3	A	602	ADP	N6-C6-N1	2.73	124.34	118.35
3	A	602	ADP	C5-C6-N6	-2.46	118.07	123.43
3	B	602	ADP	C2-N1-C6	2.33	122.77	118.77
3	B	602	ADP	PA-O5'-C5'	-2.29	108.25	121.68

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	ADP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	337/426 (79%)	-0.06	14 (4%)	40 26	30, 61, 108, 136	0
1	B	332/426 (77%)	-0.08	14 (4%)	40 26	35, 64, 110, 140	0
All	All	669/852 (78%)	-0.07	28 (4%)	40 26	30, 63, 110, 140	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	495	GLY	4.5
1	B	527	GLY	3.6
1	A	505	THR	3.4
1	B	464	SER	3.1
1	B	363	GLN	3.0
1	B	358	LEU	3.0
1	A	535	PHE	3.0
1	B	241	CYS	2.9
1	A	224	PRO	2.8
1	A	465	ARG	2.8
1	B	535	PHE	2.8
1	B	364	ASN	2.7
1	A	496	ALA	2.7
1	A	358	LEU	2.7
1	A	528	GLN	2.7
1	A	507	MET	2.6
1	B	249	PRO	2.6
1	A	504	GLN	2.6
1	B	463	SER	2.5
1	A	494	ARG	2.4
1	B	494	ARG	2.4
1	B	465	ARG	2.4
1	A	584	LEU	2.3
1	A	464	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	463	SER	2.1
1	B	366	SER	2.1
1	B	365	ALA	2.1
1	B	584	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	AF3	B	603	4/4	0.93	0.22	33,34,34,35	0
4	AF3	A	603	4/4	0.95	0.20	32,32,34,36	0
2	MG	A	601	1/1	0.96	0.08	123,123,123,123	0
3	ADP	A	602	27/27	0.97	0.06	46,53,58,66	0
3	ADP	B	602	27/27	0.97	0.07	39,47,64,72	0
2	MG	B	601	1/1	0.98	0.10	58,58,58,58	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ADP A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ADP B 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.