



wwPDB EM Validation Summary Report ⓘ

Aug 4, 2026 – 09:00 PM EDT

PDB ID : 6XKZ / pdb_00006xkz
EMDB ID : EMD-22230
Title : R. capsulatus CIII2CIV tripartite super-complex, conformation B (SC-1B)
Authors : Steimle, S.; Van Eeuwen, T.; Ozturk, Y.; Kim, H.J.; Braitbard, M.; Selamoglu, N.; Garcia, B.A.; Schneidman-Duhovny, D.; Murakami, K.; Daldal, F.
Deposited on : 2020-06-27
Resolution : 7.20 Å(reported)
Based on initial model : 6XI0

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev133
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 7.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	444 (6.70 - 7.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	191	67% 48% 46% 5%
1	R	191	79% 47% 47% 5%
2	C	437	27% 76% 19% ..
2	P	437	17% 74% 22% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	279	
3	Q	279	
4	n	532	
5	o	242	
6	p	474	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	HEC	C	502	X	-	-	-
8	HEC	D	501	X	-	-	-
8	HEC	P	502	X	-	-	-
8	HEC	Q	501	X	-	-	-
8	HEC	n	601	X	-	-	-
8	HEC	n	602	X	-	X	-
8	HEC	o	301	X	-	-	-
8	HEC	p	501	X	-	-	-
8	HEC	p	502	X	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 20406 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquinol-cytochrome c reductase iron-sulfur subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	E	181	Total	C	N	O	S	0	0
			1365	855	246	256	8		
1	R	181	Total	C	N	O	S	0	0
			1365	855	246	256	8		

- Molecule 2 is a protein called Cytochrome b.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	423	Total	C	N	O	S	0	0
			3244	2186	519	525	14		
2	P	423	Total	C	N	O	S	0	0
			3244	2186	519	525	14		

- Molecule 3 is a protein called Cytochrome c1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	235	Total	C	N	O	S	0	0
			1801	1148	301	337	15		
3	Q	235	Total	C	N	O	S	0	0
			1794	1144	300	335	15		

- Molecule 4 is a protein called Cytochrome c oxidase, Cbb3-type, subunit I.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	n	471	Total	C	N	O	S	0	0
			3711	2481	598	609	23		

- Molecule 5 is a protein called Cytochrome c oxidase, Cbb3-type, subunit II.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	o	190	Total	C	N	O	S	0	0
			1498	960	244	284	10		


- Molecule 6 is a protein called Cbb3-type cytochrome c oxidase subunit CcoP,Cytochrome c-type cyt cy.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	p	254	Total	C	N	O	S	0	0
			1902	1203	322	367	10		

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
p	467	ASP	-	expression tag	UNP Q05389
p	468	TYR	-	expression tag	UNP Q05389
p	469	LYS	-	expression tag	UNP Q05389
p	470	ASP	-	expression tag	UNP Q05389
p	471	ASP	-	expression tag	UNP Q05389
p	472	ASP	-	expression tag	UNP Q05389
p	473	ASP	-	expression tag	UNP Q05389
p	474	LYS	-	expression tag	UNP Q05389

- Molecule 7 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms			AltConf
7	E	1	Total	Fe	S	0
			4	2	2	
7	R	1	Total	Fe	S	0
			4	2	2	

- Molecule 8 is HEME C (CCD ID: HEC) (formula: $\text{C}_{34}\text{H}_{36}\text{FeN}_4\text{O}_4$).



Mol	Chain	Residues	Atoms					AltConf
8	C	1	Total 43	C 34	Fe 1	N 4	O 4	0
8	C	1	Total 43	C 34	Fe 1	N 4	O 4	0
8	D	1	Total 43	C 34	Fe 1	N 4	O 4	0
8	P	1	Total 43	C 34	Fe 1	N 4	O 4	0
8	P	1	Total 43	C 34	Fe 1	N 4	O 4	0
8	Q	1	Total 43	C 34	Fe 1	N 4	O 4	0
8	n	1	Total 43	C 34	Fe 1	N 4	O 4	0
8	n	1	Total 43	C 34	Fe 1	N 4	O 4	0
8	o	1	Total 43	C 34	Fe 1	N 4	O 4	0
8	p	1	Total 43	C 34	Fe 1	N 4	O 4	0
8	p	1	Total 43	C 34	Fe 1	N 4	O 4	0


- Molecule 9 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		AltConf
9	n	1	Total 1	Cu 1	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ubiquinol-cytochrome c reductase iron-sulfur subunit




- Molecule 1: Ubiquinol-cytochrome c reductase iron-sulfur subunit





- Molecule 2: Cytochrome b







• Molecule 2: Cytochrome b




• Molecule 3: Cytochrome c1








• Molecule 3: Cytochrome c1




• Molecule 4: Cytochrome c oxidase, Cbb3-type, subunit I






• Molecule 5: Cytochrome c oxidase, Cbb3-type, subunit II



• Molecule 6: Cbb3-type cytochrome c oxidase subunit CcoP, Cytochrome c-type cyt cy



VAL	GLU	ASP	D190
SER	GLU	GLY	V191
PRO	LYS	T251	V192
LYS	PRE	T252	Q193
ALA	ASN	T253	Y194
GLU	LYS	LEU	V195
VAL	THR	PRO	V196
PRO	CYS	SER	L196
GLY	LYS	VAL	K197
THR	ALA	ASP	G259
LYS	LYS	ASP	I198
LYS	CYS	GLU	S199
MET	HIS	ALA	G200
SER	LYS	ALA	Q201
SER	LYS	MET	P202
PHE	ILE	VAL	A203
VAL	ASP	PRO	D204
GLY	GLY	ALA	E205
GLY	LEU	S266	A206
LEU	LEU	W267	R207
PRO	PRO	ALA	A208
PRO	ASN	ALA	T209
GLU	ALA	PRO	A210
ALA	VAL	ALA	G211
ALA	GLY	ALA	Q212
ASP	PRO	ALA	Q213
ASP	ARG	GLY	W214
ARG	HIS	THR	F215
ALA	ILE	ALA	A216
ALA	TYR	ALA	D217
LEU	GLY	ALA	N218
LEU	GLY	PRO	C219
LEU	ARG	ALA	V220
ASN	ALA	GLU	S221
THR	THR	ALA	C222
THR	ALA	ALA	H223
LEU	THR	ALA	G224
PRO	LYS	PRO	E225
ASP	TYR	PRO	D226
ASP	SER	PRO	A227
LYS	LYS	PRO	K228
ASP	ALA	ALA	G229
LYS	MET	ALA	M230
ASP	THR	VAL	V231
ASP	THR	GLY	E232
LYS	THR	LEU	M233
THR	THR	ALA	G234
THR	THR	THR	A235
GLU	PRO	ARG	P236
GLU	ARG	ALA	N237
LEU	LEU	ALA	L238
ASP	ASP	THR	T239
ILE	ILE	VAL	D240
TYR	TYR	ALA	G241
LEU	LEU	VAL	I242
		GLY	W243
		ALA	L244
			Y245
			G246
			G247
			D248

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	87026	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.109	Depositor
Minimum map value	-0.044	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.018	Depositor
Map size ($\text{\AA}$)	316.80002, 316.80002, 316.80002	wwPDB
Map dimensions	240, 240, 240	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.32, 1.32, 1.32	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU, HEC, FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.35	0/1395	0.66	1/1895 (0.1%)
1	R	0.35	0/1395	0.66	1/1895 (0.1%)
2	C	0.37	0/3363	0.61	7/4617 (0.2%)
2	P	0.38	0/3363	0.62	8/4617 (0.2%)
3	D	0.28	0/1846	0.60	5/2500 (0.2%)
3	Q	0.28	0/1839	0.59	5/2492 (0.2%)
4	n	1.26	2/3836 (0.1%)	1.65	67/5232 (1.3%)
5	o	1.26	1/1535 (0.1%)	1.59	16/2092 (0.8%)
6	p	1.28	2/1948 (0.1%)	1.81	54/2660 (2.0%)
All	All	0.80	5/20520 (0.0%)	1.12	164/28000 (0.6%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	p	105	LEU	CA-C	-6.47	1.44	1.52
4	n	283	VAL	CA-CB	-6.45	1.50	1.54
6	p	188	ILE	CA-CB	-6.14	1.46	1.54
4	n	245	LYS	N-CA	-5.46	1.39	1.46
5	o	72	HIS	C-N	-5.08	1.26	1.33

The worst 5 of 164 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	p	113	GLY	N-CA-C	-11.31	99.16	112.73
4	n	282	PHE	CA-CB-CG	-10.32	103.48	113.80
2	P	222	ASN	N-CA-C	9.86	121.62	111.07
4	n	269	PHE	CA-CB-CG	-9.36	104.44	113.80
6	p	107	THR	N-CA-CB	9.24	124.03	110.26

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1365	0	1340	135	0
1	R	1365	0	1340	140	0
2	C	3244	0	3072	91	0
2	P	3244	0	3072	102	0
3	D	1801	0	1722	103	0
3	Q	1794	0	1709	73	0
4	n	3711	0	3700	1010	0
5	o	1498	0	1458	363	0
6	p	1902	0	1816	652	0
7	E	4	0	0	0	0
7	R	4	0	0	0	0
8	C	86	0	64	16	0
8	D	43	0	30	5	0
8	P	86	0	64	17	0
8	Q	43	0	30	5	0
8	n	86	0	64	34	0
8	o	43	0	30	5	0
8	p	86	0	60	45	0
9	n	1	0	0	0	0
All	All	20406	0	19571	2438	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 61.

The worst 5 of 2438 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:p:103:PRO:CA	6:p:107:THR:HG23	1.34	1.55
6:p:177:HIS:CB	6:p:183:LEU:HD23	1.35	1.55
3:D:7:HIS:H	6:p:65:LEU:CD2	1.23	1.52
6:p:177:HIS:HB3	6:p:183:LEU:CD2	1.42	1.48
3:D:6:ASP:N	6:p:65:LEU:CB	1.76	1.48

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	179/191 (94%)	141 (79%)	37 (21%)	1 (1%)	21	59
1	R	179/191 (94%)	141 (79%)	36 (20%)	2 (1%)	11	46
2	C	419/437 (96%)	352 (84%)	65 (16%)	2 (0%)	24	63
2	P	419/437 (96%)	352 (84%)	66 (16%)	1 (0%)	43	78
3	D	231/279 (83%)	198 (86%)	33 (14%)	0	100	100
3	Q	231/279 (83%)	197 (85%)	34 (15%)	0	100	100
4	n	469/532 (88%)	458 (98%)	9 (2%)	2 (0%)	30	67
5	o	186/242 (77%)	177 (95%)	9 (5%)	0	100	100
6	p	246/474 (52%)	225 (92%)	16 (6%)	5 (2%)	6	31
All	All	2559/3062 (84%)	2241 (88%)	305 (12%)	13 (0%)	26	63

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	n	289	ARG
6	p	176	ALA
1	E	99	GLU
1	R	99	GLU
6	p	182	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	142/149 (95%)	142 (100%)	0	100	100
1	R	142/149 (95%)	142 (100%)	0	100	100
2	C	300/360 (83%)	299 (100%)	1 (0%)	86	86
2	P	300/360 (83%)	300 (100%)	0	100	100
3	D	184/220 (84%)	184 (100%)	0	100	100
3	Q	182/220 (83%)	182 (100%)	0	100	100
4	n	376/421 (89%)	349 (93%)	27 (7%)	13	35
5	o	163/205 (80%)	157 (96%)	6 (4%)	30	51
6	p	192/357 (54%)	182 (95%)	10 (5%)	21	42
All	All	1981/2441 (81%)	1937 (98%)	44 (2%)	45	64

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	o	39	LEU
6	p	79	LYS
5	o	41	ASN
5	o	166	VAL
6	p	144	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
6	p	84	ASN
6	p	282	GLN
6	p	132	ASN
6	p	177	HIS
3	Q	31	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 1 is monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	HEC	p	502	6	50,50,50	1.55	8 (16%)	68,82,82	1.37	12 (17%)
8	HEC	o	301	5	50,50,50	1.56	6 (12%)	68,82,82	1.55	10 (14%)
8	HEC	P	501	2	50,50,50	1.57	9 (18%)	68,82,82	1.39	9 (13%)
8	HEC	n	602	4	50,50,50	1.91	9 (18%)	68,82,82	1.61	10 (14%)
7	FES	R	501	1	0,4,4	-	-	-	-	-
8	HEC	D	501	3	50,50,50	1.79	7 (14%)	68,82,82	1.24	2 (2%)
8	HEC	Q	501	3	50,50,50	1.79	7 (14%)	68,82,82	1.23	2 (2%)
8	HEC	n	601	4	50,50,50	1.94	11 (22%)	68,82,82	1.54	10 (14%)
8	HEC	C	501	2	50,50,50	1.57	9 (18%)	68,82,82	1.39	9 (13%)
8	HEC	P	502	2	50,50,50	1.66	9 (18%)	68,82,82	1.41	9 (13%)
8	HEC	C	502	2	50,50,50	1.66	9 (18%)	68,82,82	1.41	9 (13%)
7	FES	E	501	1	0,4,4	-	-	-	-	-
8	HEC	p	501	6	50,50,50	1.63	6 (12%)	68,82,82	1.54	8 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	HEC	p	502	6	1/1/3/7	5/14/54/54	-
8	HEC	o	301	5	1/1/3/7	6/14/54/54	-
8	HEC	P	501	2	-	8/14/54/54	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	HEC	n	602	4	1/1/3/7	6/14/54/54	-
7	FES	R	501	1	-	-	0/1/1/1
8	HEC	D	501	3	1/1/3/7	6/14/54/54	-
8	HEC	Q	501	3	1/1/3/7	6/14/54/54	-
8	HEC	n	601	4	1/1/3/7	8/14/54/54	-
8	HEC	C	501	2	-	8/14/54/54	-
8	HEC	P	502	2	1/1/3/7	9/14/54/54	-
8	HEC	C	502	2	1/1/3/7	9/14/54/54	-
7	FES	E	501	1	-	-	0/1/1/1
8	HEC	p	501	6	1/1/3/7	6/14/54/54	-

The worst 5 of 90 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	Q	501	HEC	C3D-C2D	7.74	1.53	1.36
8	D	501	HEC	C3D-C2D	7.70	1.53	1.36
8	n	601	HEC	C3D-C2D	7.67	1.53	1.36
8	n	602	HEC	C3D-C2D	7.63	1.53	1.36
8	p	501	HEC	C3D-C2D	7.32	1.52	1.36

The worst 5 of 90 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	p	501	HEC	CAC-C3C-C4C	5.89	133.01	124.91
8	o	301	HEC	CAC-C3C-C4C	5.09	131.92	124.91
8	n	601	HEC	C1D-ND-C4D	5.09	111.24	105.21
8	D	501	HEC	C1D-ND-C4D	5.03	111.16	105.21
8	Q	501	HEC	C1D-ND-C4D	5.00	111.13	105.21

5 of 9 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
8	C	502	HEC	NA
8	D	501	HEC	NA
8	P	502	HEC	NA
8	Q	501	HEC	NA
8	n	601	HEC	NA

5 of 77 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	C	501	HEC	C2C-C3C-CAC-CBC
8	P	501	HEC	C2C-C3C-CAC-CBC
8	n	602	HEC	C2B-C3B-CAB-CBB
8	n	602	HEC	C4B-C3B-CAB-CBB
8	n	601	HEC	C4C-C3C-CAC-CBC


There are no ring outliers.

11 monomers are involved in 127 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	p	502	HEC	30	0
8	o	301	HEC	5	0
8	P	501	HEC	14	0
8	n	602	HEC	22	0
8	D	501	HEC	5	0
8	Q	501	HEC	5	0
8	n	601	HEC	12	0
8	C	501	HEC	13	0
8	P	502	HEC	3	0
8	C	502	HEC	3	0
8	p	501	HEC	16	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand HEC p 502



Bond lengths



Bond angles




Torsions




Rings


Ligand HEC o 301



Bond lengths




Bond angles




Torsions




Rings




Ligand HEC n 602



Bond lengths




Bond angles




Torsions




Rings







Ligand HEC n 601



Bond lengths



Bond angles




Torsions




Rings


Ligand HEC C 501




Bond lengths



Bond angles




Torsions




Rings


Ligand HEC P 502



Bond lengths




Bond angles




Torsions



Rings





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-22230. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 120



Y Index: 120



Z Index: 120


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 122



Y Index: 112




Z Index: 86


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.018. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 196 nm^3 ; this corresponds to an approximate mass of 177 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.139 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.139 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	7.20	-	-
Author-provided FSC curve	7.18	8.86	7.29
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-22230 and PDB model 6XKZ. Per-residue inclusion information can be found in section [3](#) on page [7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.018 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.018).

9.4 Atom inclusion ⓘ



At the recommended contour level, 72% of all backbone atoms, 54% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.018) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5350	0.1220
C	0.5710	0.1350
D	0.5330	0.1100
E	0.2610	0.1020
P	0.6150	0.1410
Q	0.6450	0.1470
R	0.1600	0.0670
n	0.5930	0.1270
o	0.6080	0.1290
p	0.5260	0.0890

1.0

0.0

<0.0