



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:58 PM UTC

PDB ID : 2XXQ / pdb_00002xxq
Title : A widespread family of bacterial cell wall assembly proteins
Authors : Marles-Wright, J.; Kawai, Y.; Emmins, R.; Ishikawa, S.; Kuwano, M.; Heinz, N.; Cleverley, R.M.; Bui, N.K.; Ogasawara, N.; Lewis, R.J.; Vollmer, W.; Daniel, R.A.; Errington, J.
Deposited on : 2010-11-11
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3370 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CPS2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	373	Total	C	N	O	S	23	8	0
			2923	1836	484	595	8			

There are 15 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	97	MET	-	expression tag	UNP Q9ZII9
A	482	LYS	-	expression tag	UNP Q9ZII9
A	483	LEU	-	expression tag	UNP Q9ZII9
A	484	ALA	-	expression tag	UNP Q9ZII9
A	485	ALA	-	expression tag	UNP Q9ZII9
A	486	ALA	-	expression tag	UNP Q9ZII9
A	487	LEU	-	expression tag	UNP Q9ZII9
A	488	GLU	-	expression tag	UNP Q9ZII9
A	489	HIS	-	expression tag	UNP Q9ZII9
A	490	HIS	-	expression tag	UNP Q9ZII9
A	491	HIS	-	expression tag	UNP Q9ZII9
A	492	HIS	-	expression tag	UNP Q9ZII9
A	493	HIS	-	expression tag	UNP Q9ZII9
A	494	HIS	-	expression tag	UNP Q9ZII9
A	267	ALA	ARG	engineered mutation	UNP Q9ZII9

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is (2Z,6Z,10Z,14Z,18Z,22Z,26Z)-3,7,11,15,19,23,27,31-octamethyldotriacenta-2,6,10,14,18,22,26,30-octaen-1-yl trihydrogen diphosphate (CCD ID: ZTP) (formula: $C_{40}H_{68}O_7P_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			49	40	7	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	374	Total	O	0	0
			374	374		

- Molecule 1: CPS2A



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	73.50Å 73.50Å 163.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.64 – 1.77 37.64 – 1.77	Depositor EDS
% Data completeness (in resolution range)	99.9 (37.64-1.77) 99.9 (37.64-1.77)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.71 (at 1.77Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.170 , 0.199 0.171 , 0.199	Depositor DCC
R_{free} test set	2260 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	20.1	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3370	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZTP, ACT, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	4/2992 (0.1%)	0.81	4/4057 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	393	GLU	CD-OE2	-16.23	0.94	1.25
1	A	393	GLU	CD-OE1	14.46	1.52	1.25
1	A	214	LYS	CA-CB	-7.21	1.41	1.53
1	A	125	GLU	CG-CD	6.37	1.68	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	214	LYS	N-CA-CB	10.96	126.38	109.83
1	A	214	LYS	CA-CB-CG	7.09	128.27	114.10
1	A	393	GLU	CG-CD-OE1	-6.75	102.87	118.40
1	A	214	LYS	CB-CA-C	6.26	120.11	109.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2923	0	2932	15	0
2	A	20	0	30	2	0
3	A	4	0	3	1	0
4	A	49	0	65	6	0
5	A	374	0	0	4	0
All	All	3370	0	3030	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:264[B]:THR:HG21	5:A:2260:HOH:O	1.65	0.95
1:A:249:ILE:HG12	1:A:264[B]:THR:HG22	1.49	0.92
1:A:251:MET:SD	4:A:1489:ZTP:H18	2.32	0.69
3:A:1484:ACT:H3	5:A:2306:HOH:O	1.95	0.66
1:A:421:LEU:HD13	4:A:1489:ZTP:H13A	1.79	0.64
1:A:418[B]:MET:HE1	4:A:1489:ZTP:H1	1.85	0.57
4:A:1489:ZTP:H18B	4:A:1489:ZTP:H11A	1.91	0.52
1:A:264[B]:THR:HG23	1:A:379:GLN:HE21	1.77	0.49
1:A:389:LEU:HB3	4:A:1489:ZTP:H3B	1.94	0.49
1:A:137:THR:O	1:A:141:ASN:HB2	2.13	0.48
1:A:376:ARG:HB2	2:A:1488:EDO:H22	1.97	0.46
1:A:418[B]:MET:HE1	4:A:1489:ZTP:C1	2.46	0.45
1:A:407:ASP:OD1	5:A:2292:HOH:O	2.21	0.43
1:A:264[B]:THR:HG23	1:A:379:GLN:NE2	2.33	0.43
1:A:264[B]:THR:CG2	5:A:2260:HOH:O	2.43	0.42
1:A:258:LYS:HE3	1:A:432:TYR:CE1	2.54	0.41
1:A:376:ARG:CB	2:A:1488:EDO:H22	2.51	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/398 (96%)	374 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	336/347 (97%)	334 (99%)	2 (1%)	78	70

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	214	LYS
1	A	482	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	141	ASN
1	A	143	ASN
1	A	145	GLN
1	A	162	ASN
1	A	379	GLN
1	A	423	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	1488	-	3,3,3	0.42	0	2,2,2	0.06	0
2	EDO	A	1485	-	3,3,3	0.41	0	2,2,2	0.40	0
2	EDO	A	1483	-	3,3,3	0.39	0	2,2,2	0.52	0
3	ACT	A	1484	-	3,3,3	1.00	0	3,3,3	0.96	0
2	EDO	A	1487	-	3,3,3	0.33	0	2,2,2	0.57	0
2	EDO	A	1486	-	3,3,3	0.48	0	2,2,2	0.29	0
4	ZTP	A	1489	-	47,48,48	2.22	20 (42%)	57,61,61	1.52	12 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1488	-	-	0/1/1/1	-
2	EDO	A	1485	-	-	0/1/1/1	-
2	EDO	A	1483	-	-	0/1/1/1	-
2	EDO	A	1487	-	-	0/1/1/1	-
2	EDO	A	1486	-	-	1/1/1/1	-
4	ZTP	A	1489	-	-	12/55/55/55	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1489	ZTP	C40-C39	-3.88	1.38	1.49
4	A	1489	ZTP	C25-C24	-3.81	1.38	1.50
4	A	1489	ZTP	C20-C19	-3.57	1.39	1.50
4	A	1489	ZTP	C5-C4	-3.53	1.39	1.50
4	A	1489	ZTP	C10-C9	-3.51	1.39	1.50
4	A	1489	ZTP	C35-C34	-3.50	1.39	1.50
4	A	1489	ZTP	C30-C29	-3.47	1.40	1.50
4	A	1489	ZTP	P1-O2	3.44	1.62	1.50
4	A	1489	ZTP	C14-C12	3.43	1.41	1.33
4	A	1489	ZTP	C29-C27	3.42	1.40	1.33
4	A	1489	ZTP	C15-C14	-3.36	1.40	1.50
4	A	1489	ZTP	C19-C17	3.25	1.40	1.33
4	A	1489	ZTP	C9-C7	3.18	1.40	1.33
4	A	1489	ZTP	C24-C22	3.09	1.40	1.33
4	A	1489	ZTP	C34-C32	2.69	1.39	1.33
4	A	1489	ZTP	C39-C37	2.63	1.39	1.33
4	A	1489	ZTP	C4-C2	2.55	1.40	1.32
4	A	1489	ZTP	C11-C12	2.38	1.56	1.51
4	A	1489	ZTP	P1-O4	2.33	1.62	1.59
4	A	1489	ZTP	P2-O5	2.07	1.62	1.54

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1489	ZTP	C40-C39-C37	3.83	132.47	126.20
4	A	1489	ZTP	C33-C32-C31	3.47	121.26	115.23
4	A	1489	ZTP	C23-C22-C21	3.47	121.25	115.23
4	A	1489	ZTP	C13-C12-C14	-3.18	115.46	123.63
4	A	1489	ZTP	C38-C37-C39	-2.98	115.97	123.63
4	A	1489	ZTP	C38-C37-C36	2.74	119.99	115.23
4	A	1489	ZTP	C33-C32-C34	-2.42	117.40	123.63
4	A	1489	ZTP	C8-C7-C6	2.28	119.18	115.23
4	A	1489	ZTP	C23-C22-C24	-2.25	117.86	123.63
4	A	1489	ZTP	C18-C17-C19	-2.20	117.99	123.63
4	A	1489	ZTP	C11-C12-C14	2.16	126.03	121.17
4	A	1489	ZTP	C8-C7-C9	-2.10	118.23	123.63

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1489	ZTP	C20-C21-C22-C23

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	1489	ZTP	C20-C21-C22-C24
4	A	1489	ZTP	C29-C30-C31-C32
4	A	1489	ZTP	C19-C20-C21-C22
4	A	1489	ZTP	C24-C25-C26-C27
4	A	1489	ZTP	C14-C15-C16-C17
4	A	1489	ZTP	C9-C10-C11-C12
4	A	1489	ZTP	C10-C11-C12-C13
4	A	1489	ZTP	C10-C11-C12-C14
2	A	1486	EDO	O1-C1-C2-O2
4	A	1489	ZTP	C15-C16-C17-C18
4	A	1489	ZTP	C4-C5-C6-C7
4	A	1489	ZTP	C15-C16-C17-C19

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1488	EDO	2	0
3	A	1484	ACT	1	0
4	A	1489	ZTP	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	373/398 (93%)	-0.21	10 (2%) 56 63	7, 18, 31, 54	15 (4%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	482	LYS	4.1
1	A	215	VAL	3.5
1	A	480	GLY	3.4
1	A	481	ARG	2.9
1	A	110	SER	2.5
1	A	158	ASP	2.4
1	A	214	LYS	2.3
1	A	212	THR	2.2
1	A	340	HIS	2.1
1	A	407	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	A	1485	4/4	0.84	0.14	37,40,41,42	0
4	ZTP	A	1489	49/49	0.85	0.19	27,37,55,56	0
3	ACT	A	1484	4/4	0.88	0.14	28,29,30,31	0
2	EDO	A	1487	4/4	0.89	0.13	33,34,35,36	0
2	EDO	A	1486	4/4	0.91	0.10	29,32,34,34	0
2	EDO	A	1488	4/4	0.93	0.11	29,33,35,36	0
2	EDO	A	1483	4/4	0.95	0.08	28,29,29,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.