



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2026 – 04:01 PM JST

PDB ID : 7YJE / pdb_00007yje
Title : Crystal structure of the P450 BM3 heme domain mutant F87G/T268V/A184V/A328V in complex with N-imidazolyl-hexanoyl-L-phenylalanine and acetate ion
Authors : Dong, S.; Chen, J.; Jiang, Y.; Cong, Z.; Feng, Y.
Deposited on : 2022-07-20
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

i

X-RAY DIFFRACTION

A.

the following graphic. The table shows the number of entries on which the scores are based.

Metric

(#Entries)

 $(\# \text{Entries, resolution range}(\text{\AA}))$ R_{free}

Clashscore

Ramachandran outliers

Sidechain outliers

RSRZ outliers

the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled.

Mol

Chain

Length

Quality of chain

1

A

466

88%

•

1

B

466

90%



2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8666 atoms, of which 154 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional cytochrome P450/NADPH-P450 reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	454	Total	C	N	O	S	0	0	0
			3660	2339	622	682	17			
1	B	454	Total	C	N	O	S	0	0	0
			3660	2339	622	682	17			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	initiating methionine	UNP P14779
A	-1	GLY	-	expression tag	UNP P14779
A	87	GLY	PHE	engineered mutation	UNP P14779
A	184	VAL	ALA	engineered mutation	UNP P14779
A	268	VAL	THR	engineered mutation	UNP P14779
A	328	VAL	ALA	engineered mutation	UNP P14779
A	456	LEU	-	expression tag	UNP P14779
A	457	GLU	-	expression tag	UNP P14779
A	458	HIS	-	expression tag	UNP P14779
A	459	HIS	-	expression tag	UNP P14779
A	460	HIS	-	expression tag	UNP P14779
A	461	HIS	-	expression tag	UNP P14779
A	462	HIS	-	expression tag	UNP P14779
A	463	HIS	-	expression tag	UNP P14779
B	-2	MET	-	initiating methionine	UNP P14779
B	-1	GLY	-	expression tag	UNP P14779
B	87	GLY	PHE	engineered mutation	UNP P14779
B	184	VAL	ALA	engineered mutation	UNP P14779
B	268	VAL	THR	engineered mutation	UNP P14779
B	328	VAL	ALA	engineered mutation	UNP P14779
B	456	LEU	-	expression tag	UNP P14779
B	457	GLU	-	expression tag	UNP P14779
B	458	HIS	-	expression tag	UNP P14779
B	459	HIS	-	expression tag	UNP P14779
B	460	HIS	-	expression tag	UNP P14779

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	461	HIS	-	expression tag	UNP P14779
B	462	HIS	-	expression tag	UNP P14779
B	463	HIS	-	expression tag	UNP P14779

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 73	C 34	Fe 1	H 30	N 4	O 4	0	0
2	B	1	Total 73	C 34	Fe 1	H 30	N 4	O 4	0	0

- Molecule 3 is (2S)-2-(6-imidazol-1-ylhexanoylamino)-3-phenyl-propanoic acid (CCD ID: IC6) (formula: C₁₈H₂₃N₃O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	1
			92	36	44	6	6		
3	B	1	Total	C	H	N	O	0	1
			92	36	44	6	6		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			7	2	3	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	0
			7	2	3	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	503	Total	O	0	0
			503	503		
5	B	499	Total	O	0	0
			499	499		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase



- Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.70Å 148.51Å 64.10Å 90.00° 99.73° 90.00°	Depositor
Resolution (Å)	46.79 – 1.85 46.79 – 1.85	Depositor EDS
% Data completeness (in resolution range)	70.4 (46.79-1.85) 91.5 (46.79-1.85)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 1.86Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.146 , 0.186 0.148 , 0.187	Depositor DCC
R_{free} test set	2000 reflections (2.28%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 56.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8666	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, HEM, IC6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	1/3744 (0.0%)	0.75	1/5062 (0.0%)
1	B	0.76	0/3744	0.76	0/5062
All	All	0.77	1/7488 (0.0%)	0.76	1/10124 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	129	LYS	C-O	-5.96	1.17	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	231	ASP	CB-CA-C	5.11	119.79	112.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3660	0	3639	28	0
1	B	3660	0	3639	29	0
2	A	43	30	30	0	0
2	B	43	30	30	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	48	44	0	0	0
3	B	48	44	0	0	0
4	A	4	3	3	0	0
4	B	4	3	3	0	0
5	A	503	0	0	2	0
5	B	499	0	0	5	0
All	All	8512	154	7344	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:ARG:HD2	1:A:208:ASP:OD2	1.84	0.75
1:A:31:LYS:HE3	1:A:35:GLU:CD	2.14	0.71
1:A:377:GLU:O	1:A:380:GLU:HG2	1.91	0.69
1:B:120:VAL:HG11	1:B:302:VAL:CG2	2.24	0.67
1:B:13:GLU:H	1:B:13:GLU:CD	2.07	0.63
1:B:120:VAL:HG11	1:B:302:VAL:HG23	1.81	0.61
1:B:388:HIS:HA	1:B:391:LYS:HD3	1.81	0.61
1:A:110:GLN:NE2	1:B:386:PRO:HG2	2.17	0.58
1:A:179:ARG:HD3	1:A:204:GLN:CG	2.35	0.57
1:A:110:GLN:HE22	1:B:386:PRO:HG2	1.71	0.56
1:A:388:HIS:HA	1:A:391:LYS:HD3	1.88	0.56
1:B:195:ASP:OD1	1:B:196:PRO:HD2	2.07	0.55
1:B:306:LYS:HG2	5:B:891:HOH:O	2.06	0.55
1:B:400:CYS:HB2	2:B:501:HEM:NA	2.24	0.53
1:A:195:ASP:OD1	1:A:196:PRO:HD2	2.09	0.52
1:A:110:GLN:HE22	1:B:386:PRO:CG	2.22	0.52
1:A:381:ASN:ND2	1:A:383:SER:OG	2.43	0.52
1:A:122:ILE:HG22	1:A:148:LEU:HD22	1.92	0.51
1:A:385:ILE:O	1:B:289:LYS:HE3	2.11	0.51
1:A:367:TRP:HB2	1:A:371:VAL:HG12	1.93	0.50
1:A:171:HIS:HB2	5:A:718:HOH:O	2.12	0.50
1:B:367:TRP:HB2	1:B:371:VAL:HG12	1.94	0.49
1:A:120:VAL:HG11	1:A:302:VAL:CG1	2.43	0.48
1:B:120:VAL:HG11	1:B:302:VAL:HG21	1.94	0.48
1:B:120:VAL:CG1	1:B:302:VAL:HG23	2.44	0.48
1:A:158:PHE:CE2	1:A:219:ILE:HD13	2.49	0.48
1:B:272:LEU:HD13	1:B:322:LEU:HG	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:434:LYS:HD2	1:B:442:GLU:OE2	2.16	0.46
1:A:385:ILE:O	1:B:289:LYS:CE	2.64	0.46
1:B:167:ARG:HD2	5:B:894:HOH:O	2.15	0.46
1:A:81:PHE:HB3	1:A:209:ILE:HG12	1.98	0.46
1:A:141:VAL:HB	1:A:142:PRO:HD3	1.97	0.45
1:B:382:PRO:HD2	5:B:917:HOH:O	2.16	0.45
1:A:126:LEU:C	1:A:126:LEU:HD13	2.42	0.45
1:A:179:ARG:HD3	1:A:204:GLN:HG2	1.99	0.44
1:A:124:VAL:HG13	1:A:455:LEU:HD13	1.99	0.43
1:B:434:LYS:NZ	5:B:627:HOH:O	2.52	0.43
1:A:352:GLU:HG2	1:A:353:LEU:N	2.34	0.43
1:B:449:LYS:HA	1:B:449:LYS:HD3	1.67	0.43
1:B:186:ASN:O	1:B:190:ARG:HG3	2.19	0.42
1:B:141:VAL:HB	1:B:142:PRO:HD3	2.00	0.42
1:B:171:HIS:HB2	5:B:608:HOH:O	2.19	0.42
1:A:368:GLY:O	1:A:371:VAL:HG13	2.20	0.41
1:A:434:LYS:HD2	1:A:442:GLU:OE2	2.21	0.41
1:B:149:THR:HA	1:B:409:GLU:OE2	2.20	0.41
1:B:195:ASP:OD1	1:B:196:PRO:CD	2.67	0.41
1:A:320:GLU:HG3	1:A:374:PHE:CD1	2.56	0.41
1:A:434:LYS:NZ	5:A:618:HOH:O	2.53	0.41
1:A:120:VAL:HG11	1:A:302:VAL:HG11	2.01	0.41
1:B:124:VAL:HG13	1:B:455:LEU:HD13	2.02	0.41
1:B:280:LEU:HD22	1:B:287:LEU:HA	2.03	0.40
1:B:400:CYS:HB2	2:B:501:HEM:C4A	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	452/466 (97%)	440 (97%)	11 (2%)	1 (0%)	43	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	452/466 (97%)	439 (97%)	12 (3%)	1 (0%)	43	31
All	All	904/932 (97%)	879 (97%)	23 (2%)	2 (0%)	43	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	266	HIS
1	B	266	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	399/410 (97%)	398 (100%)	1 (0%)	86	85
1	B	399/410 (97%)	397 (100%)	2 (0%)	81	77
All	All	798/820 (97%)	795 (100%)	3 (0%)	84	82

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	128	GLN
1	B	148	LEU
1	B	380	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	204	GLN
1	A	288	GLN
1	B	70	ASN
1	B	204	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	B	501	4,1	50,50,50	1.31	6 (12%)	66,82,82	1.42	8 (12%)
4	ACT	B	503	2	3,3,3	0.67	0	3,3,3	2.37	2 (66%)
3	IC6	B	502[A]	-	25,25,25	1.63	2 (8%)	31,31,31	1.02	2 (6%)
4	ACT	A	503	2	3,3,3	1.47	0	3,3,3	1.78	1 (33%)
3	IC6	A	502[A]	-	25,25,25	1.34	1 (4%)	31,31,31	1.26	3 (9%)
3	IC6	B	502[B]	-	25,25,25	1.59	1 (4%)	31,31,31	1.07	2 (6%)
2	HEM	A	501	4,1	50,50,50	1.42	8 (16%)	66,82,82	1.35	8 (12%)
3	IC6	A	502[B]	-	25,25,25	1.65	5 (20%)	31,31,31	1.22	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	501	4,1	-	2/14/54/54	-
3	IC6	B	502[A]	-	-	1/20/20/20	0/2/2/2
3	IC6	A	502[A]	-	-	0/20/20/20	0/2/2/2
3	IC6	B	502[B]	-	-	5/20/20/20	0/2/2/2
2	HEM	A	501	4,1	-	2/14/54/54	-
3	IC6	A	502[B]	-	-	3/20/20/20	0/2/2/2

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502[A]	IC6	C11-N12	6.77	1.48	1.34
3	B	502[B]	IC6	C11-N12	6.25	1.47	1.34
3	A	502[B]	IC6	C11-N12	6.22	1.47	1.34
3	A	502[A]	IC6	C11-N12	4.67	1.44	1.34
2	B	501	HEM	FE-ND	3.55	2.05	1.94
2	A	501	HEM	CAC-C3C	3.44	1.56	1.47
2	A	501	HEM	FE-NB	2.90	2.03	1.94
2	A	501	HEM	FE-ND	2.78	2.03	1.94
2	B	501	HEM	C2A-C3A	-2.63	1.31	1.38
3	A	502[B]	IC6	C4-C3	2.58	1.41	1.35
2	A	501	HEM	FE-NC	2.56	2.03	1.95
2	A	501	HEM	FE-NA	2.54	2.03	1.95
2	B	501	HEM	FE-NB	2.49	2.02	1.94
2	A	501	HEM	CMB-C2B	2.42	1.55	1.50
3	B	502[A]	IC6	O16-C14	2.33	1.29	1.22
3	A	502[B]	IC6	O15-C14	-2.32	1.23	1.30
2	A	501	HEM	CAB-C3B	2.25	1.53	1.47
3	A	502[B]	IC6	O24-C11	-2.20	1.18	1.23
2	A	501	HEM	CMD-C2D	2.17	1.55	1.50
2	B	501	HEM	CAB-C3B	2.14	1.53	1.47
3	A	502[B]	IC6	C4-N5	-2.11	1.32	1.37
2	B	501	HEM	FE-NA	2.10	2.02	1.95
2	B	501	HEM	C1B-NB	-2.08	1.36	1.40

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	HEM	CHC-C4B-NB	4.43	129.23	124.42
2	B	501	HEM	C4D-ND-C1D	3.94	109.14	105.07
2	A	501	HEM	C4C-NC-C1C	3.42	108.69	105.35
3	A	502[A]	IC6	C09-C10-C11	-3.32	103.93	113.26
4	B	503	ACT	O-C-CH3	-3.16	110.01	122.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	HEM	C1B-NB-C4B	3.05	108.22	105.07
2	A	501	HEM	CHC-C1C-NC	2.95	127.63	124.44
3	B	502[B]	IC6	C14-C13-N12	-2.88	103.73	110.55
2	B	501	HEM	C2D-C1D-ND	-2.80	106.53	109.88
3	A	502[A]	IC6	C14-C13-N12	2.75	117.06	110.55
2	A	501	HEM	CAC-C3C-C4C	-2.72	118.32	124.90
3	A	502[A]	IC6	C17-C13-N12	-2.66	105.20	110.79
2	B	501	HEM	CBD-CAD-C3D	-2.58	105.47	112.63
4	A	503	ACT	OXT-C-O	2.56	131.50	122.05
2	B	501	HEM	C1B-NB-C4B	2.54	107.69	105.07
4	B	503	ACT	OXT-C-O	2.48	131.19	122.05
2	A	501	HEM	CHD-C1D-ND	2.47	127.10	124.42
2	A	501	HEM	CMC-C2C-C1C	-2.46	120.39	124.71
3	B	502[A]	IC6	C14-C13-N12	2.35	116.12	110.55
2	A	501	HEM	C4A-NA-C1A	2.25	107.56	105.35
3	A	502[B]	IC6	C17-C18-C23	-2.23	116.47	120.91
2	B	501	HEM	CAC-C3C-C4C	-2.20	119.58	124.90
3	A	502[B]	IC6	C4-C3-N2	-2.17	105.76	109.92
2	A	501	HEM	CBA-CAA-C2A	-2.14	106.69	112.63
3	B	502[B]	IC6	O15-C14-C13	2.10	120.39	113.40
2	B	501	HEM	CHC-C1C-NC	-2.09	122.18	124.44
3	B	502[A]	IC6	O15-C14-C13	2.08	120.32	113.40
3	A	502[B]	IC6	C17-C13-C14	-2.05	105.50	110.42
2	B	501	HEM	O1D-CGD-CBD	-2.03	116.56	123.08

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	502[B]	IC6	C8-C09-C10-C11
3	A	502[B]	IC6	C09-C10-C11-N12
3	A	502[B]	IC6	C09-C10-C11-O24
3	B	502[B]	IC6	C09-C10-C11-N12
3	B	502[B]	IC6	C09-C10-C11-O24
3	B	502[A]	IC6	C10-C09-C8-C7
3	B	502[B]	IC6	C6-C7-C8-C09
3	A	502[B]	IC6	C6-C7-C8-C09
2	A	501	HEM	CAD-CBD-CGD-O1D
2	B	501	HEM	CAD-CBD-CGD-O1D
2	A	501	HEM	CAD-CBD-CGD-O2D
2	B	501	HEM	CAD-CBD-CGD-O2D
3	B	502[B]	IC6	C14-C13-N12-C11


There are no ring outliers.


1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.













Ligand IC6 A 502 (A)



Ligand IC6 B 502 (B)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	454/466 (97%)	-0.55	2 (0%) 88 91	11, 20, 42, 62	0
1	B	454/466 (97%)	-0.54	5 (1%) 78 81	13, 21, 42, 78	0
All	All	908/932 (97%)	-0.55	7 (0%) 82 86	11, 21, 42, 78	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	383	SER	4.2
1	B	2	ILE	3.8
1	A	110	GLN	3.2
1	B	382	PRO	2.8
1	A	381	ASN	2.5
1	B	384	ALA	2.3
1	B	369	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	IC6	A	502[A]	24/24	0.90	0.09	1,16,25,25	46
3	IC6	A	502[B]	24/24	0.90	0.09	1,16,24,28	46
3	IC6	B	502[A]	24/24	0.91	0.09	3,16,22,23	46
3	IC6	B	502[B]	24/24	0.91	0.09	1,14,20,23	46
4	ACT	B	503	4/4	0.91	0.12	27,34,40,40	0
4	ACT	A	503	4/4	0.92	0.09	26,32,32,34	0
2	HEM	A	501	43/43	0.99	0.04	12,16,23,24	0
2	HEM	B	501	43/43	0.99	0.03	11,15,20,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around IC6 A 502 (A):

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around IC6 A 502 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around IC6 B 502 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around IC6 B 502 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around ACT B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ACT A 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.