



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2026 – 02:30 PM EDT

PDB ID : 2Z47 / pdb_00002z47
Title : The Y66L mutant of tetraheme cytochrome c3 from *Desulfovibrio Vulgaris*
Miyazaki F
Authors : Higuchi, Y.; Komori, H.
Deposited on : 2007-06-13
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	107	 4% 93% 7%
1	B	107	 4% 92% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HEC	A	1001	X	-	-	-
2	HEC	A	1002	X	-	-	-
2	HEC	A	1003	X	-	-	-
2	HEC	A	1004	X	-	-	-
2	HEC	B	3001	X	-	-	-
2	HEC	B	3002	X	-	-	-
2	HEC	B	3003	X	-	-	-
2	HEC	B	3004	X	-	-	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2169 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cytochrome c3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	107	Total	C	N	O	S	0	0	0
			797	486	150	150	11			
1	B	107	Total	C	N	O	S	0	0	0
			797	486	150	150	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	66	LEU	TYR	engineered mutation	UNP P00132
B	2066	LEU	TYR	engineered mutation	UNP P00132

- Molecule 2 is HEME C (CCD ID: HEC) (formula: $C_{34}H_{36}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	97	Total 97	O 97	0	0
3	B	134	Total 134	O 134	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cytochrome c3



- Molecule 1: Cytochrome c3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	34.42Å 69.56Å 53.75Å 90.00° 92.73° 90.00°	Depositor
Resolution (Å)	30.83 – 1.60 30.83 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.5 (30.83-1.60) 99.7 (30.83-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.94 (at 1.60Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.184 , 0.216 0.181 , 0.211	Depositor DCC
R_{free} test set	3330 reflections (9.99%)	wwPDB-VP
Wilson B-factor (Å ²)	13.4	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.064 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2169	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/813	0.90	3/1082 (0.3%)
1	B	0.30	0/813	0.89	3/1082 (0.3%)
All	All	0.29	0/1626	0.90	6/2164 (0.3%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	71	ASP	N-CA-C	8.25	121.69	110.55
1	B	2071	ASP	N-CA-C	7.75	121.02	110.55
1	A	70	HIS	N-CA-C	6.53	121.91	113.88
1	B	2012	ASP	N-CA-C	5.54	117.09	110.44
1	B	2070	HIS	N-CA-C	5.31	121.16	114.31
1	A	12	ASP	N-CA-C	5.25	116.73	110.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	797	0	782	3	0
1	B	797	0	779	4	0
2	A	172	0	120	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	172	0	120	3	0
3	A	97	0	0	0	0
3	B	134	0	0	0	0
All	All	2169	0	1801	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2074:THR:HB	2:B:3002:HEC:HAA1	1.91	0.53
2:B:3002:HEC:HBC3	2:B:3002:HEC:HMC1	1.95	0.48
2:A:1001:HEC:HMC1	2:A:1001:HEC:HBC3	1.99	0.44
1:A:53:ASP:OD1	1:A:63:LYS:HE3	2.18	0.43
1:A:74:THR:HB	2:A:1002:HEC:HAA1	2.01	0.42
1:B:2009:LEU:HD23	1:B:2009:LEU:C	2.45	0.42
2:A:1003:HEC:HBB3	2:A:1003:HEC:HMB1	2.03	0.41
1:A:9:LEU:HD23	1:A:9:LEU:C	2.45	0.41
1:B:2079:CYS:HA	2:B:3003:HEC:CHC	2.51	0.40
1:B:2054:ASN:ND2	1:B:2059:ASP:OD2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	105/107 (98%)	102 (97%)	2 (2%)	1 (1%)	12 3
1	B	105/107 (98%)	104 (99%)	1 (1%)	0	100 100
All	All	210/214 (98%)	206 (98%)	3 (1%)	1 (0%)	24 10

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/85 (100%)	85 (100%)	0	100	100
1	B	85/85 (100%)	84 (99%)	1 (1%)	63	43
All	All	170/170 (100%)	169 (99%)	1 (1%)	78	66

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2013	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEC	B	3004	1	50,50,50	2.58	21 (42%)	68,82,82	1.23	6 (8%)
2	HEC	A	1004	1	50,50,50	2.59	22 (44%)	68,82,82	1.20	7 (10%)
2	HEC	A	1002	1	50,50,50	2.52	21 (42%)	68,82,82	1.15	4 (5%)
2	HEC	B	3003	1	50,50,50	2.63	21 (42%)	68,82,82	1.30	9 (13%)
2	HEC	B	3002	1	50,50,50	2.57	21 (42%)	68,82,82	1.19	6 (8%)
2	HEC	A	1001	1	50,50,50	2.63	21 (42%)	68,82,82	1.26	6 (8%)
2	HEC	A	1003	1	50,50,50	2.66	21 (42%)	68,82,82	1.30	7 (10%)
2	HEC	B	3001	1	50,50,50	2.54	21 (42%)	68,82,82	1.22	4 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	B	3004	1	1/1/3/7	7/14/54/54	-
2	HEC	A	1004	1	1/1/3/7	9/14/54/54	-
2	HEC	A	1002	1	1/1/3/7	4/14/54/54	-
2	HEC	B	3003	1	1/1/3/7	7/14/54/54	-
2	HEC	B	3002	1	1/1/3/7	6/14/54/54	-
2	HEC	A	1001	1	1/1/3/7	8/14/54/54	-
2	HEC	A	1003	1	1/1/3/7	7/14/54/54	-
2	HEC	B	3001	1	1/1/3/7	6/14/54/54	-

All (169) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1002	HEC	FE-NB	5.55	2.12	1.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3002	HEC	FE-NB	5.49	2.11	1.94
2	A	1001	HEC	FE-NC	5.46	2.13	1.95
2	B	3003	HEC	FE-NB	5.42	2.11	1.94
2	A	1004	HEC	FE-NB	5.39	2.11	1.94
2	A	1001	HEC	FE-NB	5.27	2.11	1.94
2	A	1001	HEC	FE-ND	5.16	2.10	1.94
2	A	1003	HEC	FE-NA	5.09	2.12	1.95
2	B	3004	HEC	FE-NB	5.03	2.10	1.94
2	B	3001	HEC	FE-ND	5.02	2.10	1.94
2	A	1002	HEC	FE-ND	4.94	2.10	1.94
2	A	1003	HEC	FE-NB	4.94	2.10	1.94
2	B	3003	HEC	FE-NA	4.90	2.11	1.95
2	A	1003	HEC	FE-ND	4.89	2.10	1.94
2	B	3004	HEC	C1D-ND	-4.87	1.31	1.40
2	B	3002	HEC	FE-ND	4.84	2.09	1.94
2	A	1004	HEC	FE-NC	4.73	2.10	1.95
2	A	1004	HEC	FE-ND	4.72	2.09	1.94
2	B	3002	HEC	FE-NC	4.72	2.10	1.95
2	A	1004	HEC	FE-NA	4.70	2.10	1.95
2	A	1003	HEC	FE-NC	4.68	2.10	1.95
2	B	3001	HEC	FE-NB	4.67	2.09	1.94
2	B	3004	HEC	FE-NC	4.66	2.10	1.95
2	B	3003	HEC	C1D-ND	-4.65	1.32	1.40
2	B	3002	HEC	FE-NA	4.63	2.10	1.95
2	B	3003	HEC	FE-NC	4.62	2.10	1.95
2	B	3003	HEC	FE-ND	4.59	2.09	1.94
2	A	1002	HEC	FE-NC	4.57	2.10	1.95
2	B	3004	HEC	FE-NA	4.55	2.10	1.95
2	B	3004	HEC	FE-ND	4.47	2.08	1.94
2	B	3001	HEC	CAB-C3B	-4.45	1.39	1.51
2	B	3001	HEC	FE-NC	4.41	2.09	1.95
2	A	1003	HEC	C1D-ND	-4.37	1.32	1.40
2	A	1002	HEC	CAC-C3C	-4.37	1.39	1.51
2	A	1003	HEC	C4B-NB	-4.36	1.32	1.40
2	B	3001	HEC	FE-NA	4.35	2.09	1.95
2	A	1002	HEC	C1D-ND	-4.34	1.32	1.40
2	B	3003	HEC	C4D-C3D	-4.33	1.37	1.45
2	A	1004	HEC	CAB-C3B	-4.33	1.40	1.51
2	A	1002	HEC	FE-NA	4.32	2.09	1.95
2	A	1001	HEC	CAB-C3B	-4.32	1.40	1.51
2	A	1001	HEC	FE-NA	4.28	2.09	1.95
2	B	3001	HEC	C4B-NB	-4.26	1.32	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3004	HEC	CAB-C3B	-4.26	1.40	1.51
2	B	3003	HEC	C4B-NB	-4.24	1.32	1.40
2	A	1004	HEC	CAC-C3C	-4.23	1.40	1.51
2	A	1004	HEC	C4D-C3D	-4.22	1.37	1.45
2	B	3002	HEC	C1D-ND	-4.22	1.32	1.40
2	A	1003	HEC	CAB-C3B	-4.22	1.40	1.51
2	B	3002	HEC	CAC-C3C	-4.20	1.40	1.51
2	B	3001	HEC	C1D-ND	-4.20	1.33	1.40
2	A	1003	HEC	C4D-C3D	-4.20	1.37	1.45
2	A	1003	HEC	CAC-C3C	-4.18	1.40	1.51
2	A	1001	HEC	C1D-ND	-4.12	1.33	1.40
2	A	1002	HEC	CAB-C3B	-4.12	1.40	1.51
2	A	1001	HEC	CAC-C3C	-4.12	1.40	1.51
2	B	3002	HEC	CAB-C3B	-4.09	1.40	1.51
2	A	1003	HEC	C1A-C2A	-4.09	1.38	1.45
2	B	3003	HEC	CAC-C3C	-4.04	1.40	1.51
2	B	3003	HEC	CAB-C3B	-4.03	1.40	1.51
2	B	3001	HEC	CAC-C3C	-4.00	1.40	1.51
2	B	3004	HEC	CAC-C3C	-3.99	1.40	1.51
2	B	3004	HEC	C4D-C3D	-3.97	1.38	1.45
2	A	1004	HEC	C4A-NA	-3.96	1.32	1.39
2	A	1001	HEC	C4D-C3D	-3.94	1.38	1.45
2	A	1001	HEC	C4A-C3A	-3.93	1.37	1.45
2	B	3002	HEC	C4D-C3D	-3.92	1.38	1.45
2	B	3004	HEC	C4A-C3A	-3.90	1.37	1.45
2	A	1001	HEC	C4B-NB	-3.88	1.33	1.40
2	A	1004	HEC	C4B-NB	-3.86	1.33	1.40
2	B	3003	HEC	C1C-NC	-3.82	1.32	1.39
2	A	1003	HEC	C4C-NC	-3.79	1.32	1.39
2	B	3001	HEC	C4D-C3D	-3.78	1.38	1.45
2	B	3004	HEC	C4B-NB	-3.78	1.33	1.40
2	B	3002	HEC	C4B-NB	-3.77	1.33	1.40
2	B	3003	HEC	C1A-C2A	-3.72	1.38	1.45
2	B	3003	HEC	C1A-NA	-3.70	1.32	1.39
2	B	3003	HEC	C4A-NA	-3.69	1.32	1.39
2	B	3001	HEC	C1C-NC	-3.69	1.32	1.39
2	B	3004	HEC	C1A-NA	-3.67	1.32	1.39
2	A	1001	HEC	C1A-NA	-3.66	1.32	1.39
2	B	3002	HEC	C4B-C3B	-3.66	1.38	1.45
2	A	1004	HEC	C1D-ND	-3.65	1.33	1.40
2	A	1003	HEC	C1A-NA	-3.65	1.32	1.39
2	A	1004	HEC	C1A-C2A	-3.64	1.38	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3004	HEC	C4C-NC	-3.61	1.32	1.39
2	B	3002	HEC	C4A-C3A	-3.59	1.38	1.45
2	B	3001	HEC	C4B-C3B	-3.57	1.39	1.45
2	A	1003	HEC	C4B-C3B	-3.53	1.39	1.45
2	A	1002	HEC	C4A-C3A	-3.53	1.38	1.45
2	A	1002	HEC	C4B-NB	-3.52	1.34	1.40
2	A	1003	HEC	C4D-ND	-3.52	1.31	1.38
2	A	1002	HEC	C4D-C3D	-3.51	1.39	1.45
2	B	3001	HEC	C4A-C3A	-3.50	1.38	1.45
2	A	1001	HEC	C4A-NA	-3.48	1.33	1.39
2	A	1003	HEC	C1C-NC	-3.48	1.33	1.39
2	B	3004	HEC	C4B-C3B	-3.45	1.39	1.45
2	B	3003	HEC	C4B-C3B	-3.45	1.39	1.45
2	B	3003	HEC	C4D-ND	-3.44	1.32	1.38
2	A	1003	HEC	C4A-NA	-3.42	1.33	1.39
2	B	3002	HEC	C4A-NA	-3.41	1.33	1.39
2	A	1001	HEC	C4B-C3B	-3.40	1.39	1.45
2	A	1004	HEC	C4D-ND	-3.39	1.32	1.38
2	A	1004	HEC	C4B-C3B	-3.36	1.39	1.45
2	B	3003	HEC	C4C-NC	-3.35	1.33	1.39
2	A	1002	HEC	C1C-NC	-3.34	1.33	1.39
2	A	1001	HEC	C1A-C2A	-3.34	1.39	1.45
2	A	1003	HEC	C4A-C3A	-3.34	1.38	1.45
2	B	3004	HEC	C4A-NA	-3.32	1.33	1.39
2	A	1002	HEC	C4C-NC	-3.31	1.33	1.39
2	A	1001	HEC	C1D-C2D	-3.31	1.37	1.44
2	B	3001	HEC	C1A-NA	-3.31	1.33	1.39
2	B	3004	HEC	C1D-C2D	-3.30	1.37	1.44
2	B	3002	HEC	C4C-NC	-3.29	1.33	1.39
2	B	3002	HEC	C1C-NC	-3.27	1.33	1.39
2	A	1004	HEC	C4C-NC	-3.26	1.33	1.39
2	A	1004	HEC	C1A-NA	-3.26	1.33	1.39
2	B	3001	HEC	C1A-C2A	-3.25	1.39	1.45
2	B	3004	HEC	C4D-ND	-3.23	1.32	1.38
2	A	1002	HEC	C1A-NA	-3.21	1.33	1.39
2	B	3001	HEC	C4A-NA	-3.21	1.33	1.39
2	B	3002	HEC	C1A-NA	-3.20	1.33	1.39
2	A	1002	HEC	C4D-ND	-3.19	1.32	1.38
2	A	1002	HEC	C4B-C3B	-3.18	1.39	1.45
2	B	3002	HEC	C1A-C2A	-3.17	1.39	1.45
2	A	1001	HEC	C4C-NC	-3.14	1.33	1.39
2	B	3001	HEC	C4C-NC	-3.14	1.33	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3001	HEC	C1D-C2D	-3.12	1.38	1.44
2	A	1004	HEC	C4A-C3A	-3.08	1.39	1.45
2	A	1004	HEC	C1C-NC	-3.07	1.33	1.39
2	B	3002	HEC	C4D-ND	-3.07	1.32	1.38
2	A	1002	HEC	C4A-NA	-3.03	1.33	1.39
2	B	3004	HEC	C1B-NB	-3.03	1.32	1.38
2	A	1001	HEC	C4D-ND	-2.99	1.33	1.38
2	B	3004	HEC	C1C-NC	-2.99	1.34	1.39
2	B	3003	HEC	C4A-C3A	-2.99	1.39	1.45
2	B	3001	HEC	C1B-C2B	-2.99	1.38	1.44
2	A	1001	HEC	C1C-NC	-2.98	1.34	1.39
2	B	3001	HEC	C4D-ND	-2.97	1.33	1.38
2	A	1004	HEC	C1B-NB	-2.96	1.33	1.38
2	B	3004	HEC	C1B-C2B	-2.95	1.38	1.44
2	A	1003	HEC	C1D-C2D	-2.93	1.38	1.44
2	A	1002	HEC	C1A-C2A	-2.90	1.40	1.45
2	B	3002	HEC	C1D-C2D	-2.88	1.38	1.44
2	B	3003	HEC	C1D-C2D	-2.83	1.38	1.44
2	A	1003	HEC	C4C-C3C	-2.82	1.38	1.44
2	A	1004	HEC	C1B-C2B	-2.80	1.38	1.44
2	B	3004	HEC	C1A-C2A	-2.79	1.40	1.45
2	A	1002	HEC	C1D-C2D	-2.78	1.38	1.44
2	B	3003	HEC	C1B-NB	-2.77	1.33	1.38
2	A	1003	HEC	C1B-NB	-2.75	1.33	1.38
2	A	1001	HEC	C1B-C2B	-2.70	1.39	1.44
2	A	1001	HEC	C1B-NB	-2.67	1.33	1.38
2	A	1004	HEC	C1D-C2D	-2.67	1.39	1.44
2	B	3002	HEC	C1B-C2B	-2.64	1.39	1.44
2	B	3003	HEC	C4C-C3C	-2.62	1.39	1.44
2	B	3002	HEC	C4C-C3C	-2.62	1.39	1.44
2	A	1003	HEC	C1B-C2B	-2.60	1.39	1.44
2	B	3001	HEC	C1B-NB	-2.59	1.33	1.38
2	A	1001	HEC	C4C-C3C	-2.59	1.39	1.44
2	A	1002	HEC	C4C-C3C	-2.57	1.39	1.44
2	B	3001	HEC	C4C-C3C	-2.56	1.39	1.44
2	A	1004	HEC	C4C-C3C	-2.56	1.39	1.44
2	B	3003	HEC	C1B-C2B	-2.53	1.39	1.44
2	A	1002	HEC	C1B-C2B	-2.48	1.39	1.44
2	A	1002	HEC	C1B-NB	-2.45	1.34	1.38
2	B	3004	HEC	C4C-C3C	-2.41	1.39	1.44
2	B	3002	HEC	C1B-NB	-2.38	1.34	1.38
2	A	1004	HEC	C1C-C2C	-2.06	1.38	1.43

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1003	HEC	CBC-CAC-C3C	4.28	124.02	112.42
2	A	1001	HEC	CBB-CAB-C3B	4.05	123.39	112.42
2	A	1001	HEC	CBC-CAC-C3C	3.94	123.11	112.42
2	B	3003	HEC	CBC-CAC-C3C	3.87	122.91	112.42
2	B	3001	HEC	CBB-CAB-C3B	3.83	122.81	112.42
2	B	3004	HEC	CBC-CAC-C3C	3.81	122.74	112.42
2	B	3001	HEC	CBC-CAC-C3C	3.71	122.48	112.42
2	A	1003	HEC	CBB-CAB-C3B	3.42	121.70	112.42
2	A	1004	HEC	CBC-CAC-C3C	3.42	121.69	112.42
2	B	3001	HEC	C4B-NB-C1B	3.37	109.19	105.21
2	B	3002	HEC	CBC-CAC-C3C	3.18	121.03	112.42
2	B	3004	HEC	CAA-C2A-C1A	3.18	131.31	124.85
2	B	3002	HEC	C1D-ND-C4D	3.16	108.95	105.21
2	B	3003	HEC	CBB-CAB-C3B	3.15	120.95	112.42
2	A	1003	HEC	C1D-ND-C4D	3.12	108.91	105.21
2	B	3004	HEC	C1D-ND-C4D	3.08	108.85	105.21
2	A	1003	HEC	C4B-NB-C1B	3.04	108.80	105.21
2	B	3002	HEC	CBB-CAB-C3B	3.02	120.60	112.42
2	A	1002	HEC	C1D-ND-C4D	3.01	108.77	105.21
2	B	3003	HEC	C1D-ND-C4D	2.96	108.72	105.21
2	A	1001	HEC	C4B-NB-C1B	2.95	108.70	105.21
2	B	3004	HEC	C4B-NB-C1B	2.92	108.67	105.21
2	A	1004	HEC	C1D-ND-C4D	2.85	108.58	105.21
2	A	1004	HEC	C4B-NB-C1B	2.78	108.50	105.21
2	A	1002	HEC	CBC-CAC-C3C	2.78	119.97	112.42
2	B	3004	HEC	CAA-C2A-C3A	-2.74	122.74	127.87
2	A	1002	HEC	C4B-NB-C1B	2.72	108.43	105.21
2	A	1001	HEC	C1D-ND-C4D	2.62	108.31	105.21
2	A	1001	HEC	C4A-NA-C1A	2.60	110.06	105.82
2	A	1002	HEC	CBB-CAB-C3B	2.58	119.42	112.42
2	B	3003	HEC	CMC-C2C-C3C	2.55	131.03	125.62
2	A	1004	HEC	CBB-CAB-C3B	2.54	119.31	112.42
2	B	3003	HEC	CAD-C3D-C2D	2.54	132.62	127.87
2	B	3001	HEC	C1D-ND-C4D	2.53	108.20	105.21
2	A	1003	HEC	CMC-C2C-C3C	2.50	130.92	125.62
2	A	1001	HEC	CAA-C2A-C1A	2.40	129.73	124.85
2	B	3003	HEC	CAD-C3D-C4D	-2.38	120.56	124.70
2	B	3003	HEC	C4B-NB-C1B	2.36	108.00	105.21
2	B	3003	HEC	C1A-CHA-C4D	2.27	130.84	126.02
2	B	3002	HEC	C4B-NB-C1B	2.24	107.86	105.21
2	A	1003	HEC	CAD-C3D-C2D	2.21	132.00	127.87
2	A	1004	HEC	C4A-CHB-C1B	2.17	130.63	126.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3002	HEC	C4A-NA-C1A	2.15	109.33	105.82
2	B	3004	HEC	C4A-NA-C1A	2.14	109.31	105.82
2	B	3003	HEC	C1D-C2D-C3D	-2.10	104.78	106.98
2	A	1004	HEC	C4A-NA-C1A	2.09	109.22	105.82
2	A	1004	HEC	C1D-C2D-C3D	-2.05	104.82	106.98
2	B	3002	HEC	CAA-C2A-C1A	2.02	128.96	124.85
2	A	1003	HEC	C4A-CHB-C1B	2.01	130.30	126.02

All (8) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	1004	HEC	NA
2	A	1002	HEC	NA
2	A	1001	HEC	NA
2	A	1003	HEC	NA
2	B	3004	HEC	NA
2	B	3002	HEC	NA
2	B	3001	HEC	NA
2	B	3003	HEC	NA

All (54) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1001	HEC	C2C-C3C-CAC-CBC
2	B	3004	HEC	C2C-C3C-CAC-CBC
2	A	1001	HEC	C4B-C3B-CAB-CBB
2	B	3001	HEC	C2C-C3C-CAC-CBC
2	A	1003	HEC	C4B-C3B-CAB-CBB
2	B	3001	HEC	C4B-C3B-CAB-CBB
2	B	3003	HEC	C4B-C3B-CAB-CBB
2	A	1001	HEC	C2B-C3B-CAB-CBB
2	A	1003	HEC	C2B-C3B-CAB-CBB
2	B	3001	HEC	C2B-C3B-CAB-CBB
2	A	1001	HEC	C4C-C3C-CAC-CBC
2	A	1003	HEC	C4C-C3C-CAC-CBC
2	B	3003	HEC	C4C-C3C-CAC-CBC
2	B	3003	HEC	C2B-C3B-CAB-CBB
2	A	1003	HEC	C2C-C3C-CAC-CBC
2	B	3003	HEC	C2C-C3C-CAC-CBC
2	B	3004	HEC	C4C-C3C-CAC-CBC
2	B	3001	HEC	C4C-C3C-CAC-CBC
2	B	3002	HEC	C2B-C3B-CAB-CBB

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	1004	HEC	C2C-C3C-CAC-CBC
2	A	1002	HEC	C2B-C3B-CAB-CBB
2	B	3002	HEC	C4B-C3B-CAB-CBB
2	A	1002	HEC	C2C-C3C-CAC-CBC
2	A	1004	HEC	C4C-C3C-CAC-CBC
2	B	3002	HEC	C2C-C3C-CAC-CBC
2	A	1002	HEC	C4B-C3B-CAB-CBB
2	A	1004	HEC	C1A-C2A-CAA-CBA
2	A	1004	HEC	C3A-C2A-CAA-CBA
2	A	1002	HEC	C4C-C3C-CAC-CBC
2	B	3002	HEC	C4C-C3C-CAC-CBC
2	B	3001	HEC	CAA-CBA-CGA-O2A
2	B	3001	HEC	CAA-CBA-CGA-O1A
2	B	3004	HEC	C2A-CAA-CBA-CGA
2	B	3003	HEC	CAA-CBA-CGA-O2A
2	A	1001	HEC	CAA-CBA-CGA-O1A
2	A	1004	HEC	C2B-C3B-CAB-CBB
2	B	3004	HEC	CAA-CBA-CGA-O1A
2	A	1001	HEC	CAA-CBA-CGA-O2A
2	B	3003	HEC	CAA-CBA-CGA-O1A
2	B	3004	HEC	CAA-CBA-CGA-O2A
2	B	3004	HEC	CAD-CBD-CGD-O2D
2	A	1003	HEC	CAA-CBA-CGA-O2A
2	A	1004	HEC	CAA-CBA-CGA-O2A
2	B	3002	HEC	CAD-CBD-CGD-O2D
2	A	1003	HEC	CAA-CBA-CGA-O1A
2	B	3004	HEC	CAD-CBD-CGD-O1D
2	A	1004	HEC	CAD-CBD-CGD-O2D
2	A	1001	HEC	CAD-CBD-CGD-O2D
2	B	3002	HEC	CAD-CBD-CGD-O1D
2	A	1003	HEC	C3D-CAD-CBD-CGD
2	A	1004	HEC	CAA-CBA-CGA-O1A
2	A	1004	HEC	CAD-CBD-CGD-O1D
2	A	1001	HEC	CAD-CBD-CGD-O1D
2	B	3003	HEC	C3D-CAD-CBD-CGD

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1002	HEC	1	0
2	B	3003	HEC	1	0


Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3002	HEC	2	0
2	A	1001	HEC	1	0
2	A	1003	HEC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand HEC B 3004



Bond lengths



Bond angles




Torsions




Rings


Ligand HEC A 1004



Bond lengths



Bond angles




Torsions




Rings


Ligand HEC A 1002



Bond lengths



Bond angles




Torsions




Rings


Ligand HEC B 3003



Bond lengths



Bond angles




Torsions




Rings


Ligand HEC B 3002



Bond lengths



Bond angles




Torsions




Rings


Ligand HEC A 1001



Bond lengths



Bond angles




Torsions




Rings


Ligand HEC A 1003



Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	107/107 (100%)	0.41	4 (3%) 45 47	10, 17, 26, 37	0
1	B	107/107 (100%)	0.10	4 (3%) 45 47	10, 15, 22, 31	0
All	All	214/214 (100%)	0.26	8 (3%) 45 47	10, 16, 25, 37	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	ALA	8.0
1	A	2	PRO	5.0
1	B	2001	ALA	3.9
1	B	2107	SER	3.6
1	B	2056	ASP	3.2
1	A	107	SER	2.6
1	B	2057	LYS	2.3
1	A	26	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	HEC	A	1002	43/43	0.97	0.08	9,12,20,26	0
2	HEC	A	1001	43/43	0.97	0.07	11,14,17,21	0
2	HEC	A	1003	43/43	0.97	0.07	10,13,21,26	0
2	HEC	A	1004	43/43	0.98	0.06	8,10,16,22	0
2	HEC	B	3004	43/43	0.98	0.06	7,9,15,20	0
2	HEC	B	3002	43/43	0.98	0.07	8,10,20,23	0
2	HEC	B	3001	43/43	0.98	0.07	8,11,17,20	0
2	HEC	B	3003	43/43	0.98	0.06	8,10,17,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HEC A 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 1003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 1004:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC B 3004:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC B 3002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC B 3001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEC B 3003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.