



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 12:01 PM UTC

PDB ID : 1ZLK / pdb_00001zlk
Title : Crystal Structure of the Mycobacterium tuberculosis Hypoxic Response Regulator DosR C-terminal Domain-DNA Complex
Authors : Wisedchaisri, G.; Wu, M.; Rice, A.E.; Roberts, D.M.; Sherman, D.R.; Hol, W.G.J.
Deposited on : 2005-05-06
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	43	
2	D	43	
3	A	95	
3	B	95	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2018 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*GP*GP*CP*CP*CP*GP*CP*GP*CP*TP*TP*T
P*GP*GP*GP*GP*AP*CP*TP*AP*AP*AP*GP*TP*CP*CP*CP*TP*AP*AP*CP*CP*
CP*TP*GP*GP*CP*CP*AP*CP*GP*AP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	24	Total	C	N	O	P	0	0	0
			492	234	90	144	24			

- Molecule 2 is a DNA chain called 5'-D(*CP*GP*TP*GP*GP*CP*CP*AP*GP*GP*GP*T
P*TP*AP*GP*GP*GP*AP*CP*TP*TP*TP*AP*GP*TP*CP*CP*CP*CP*AP*AP*AP*
GP*CP*GP*CP*GP*GP*GP*CP*CP*AP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	25	Total	C	N	O	P	0	0	0
			516	245	97	149	25			

- Molecule 3 is a protein called Dormancy Survival Regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	65	Total	C	N	O	S	0	0	0
			505	317	90	96	2			
3	B	65	Total	C	N	O	S	0	0	0
			505	317	90	96	2			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	123	MET	-	cloning artifact	GB 15610269
A	124	GLY	-	cloning artifact	GB 15610269
A	125	SER	-	cloning artifact	GB 15610269
A	126	SER	-	cloning artifact	GB 15610269
A	127	HIS	-	expression tag	GB 15610269
A	128	HIS	-	expression tag	GB 15610269
A	129	HIS	-	expression tag	GB 15610269
A	130	HIS	-	expression tag	GB 15610269

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	HIS	-	expression tag	GB 15610269
A	132	HIS	-	expression tag	GB 15610269
A	133	SER	-	cloning artifact	GB 15610269
A	134	SER	-	cloning artifact	GB 15610269
A	135	GLY	-	cloning artifact	GB 15610269
A	136	LEU	-	cloning artifact	GB 15610269
A	137	VAL	-	cloning artifact	GB 15610269
A	138	PRO	-	cloning artifact	GB 15610269
A	139	ARG	-	cloning artifact	GB 15610269
A	140	GLY	-	cloning artifact	GB 15610269
A	141	SER	-	cloning artifact	GB 15610269
A	142	HIS	-	cloning artifact	GB 15610269
A	143	MET	-	cloning artifact	GB 15610269
B	123	MET	-	cloning artifact	GB 15610269
B	124	GLY	-	cloning artifact	GB 15610269
B	125	SER	-	cloning artifact	GB 15610269
B	126	SER	-	cloning artifact	GB 15610269
B	127	HIS	-	expression tag	GB 15610269
B	128	HIS	-	expression tag	GB 15610269
B	129	HIS	-	expression tag	GB 15610269
B	130	HIS	-	expression tag	GB 15610269
B	131	HIS	-	expression tag	GB 15610269
B	132	HIS	-	expression tag	GB 15610269
B	133	SER	-	cloning artifact	GB 15610269
B	134	SER	-	cloning artifact	GB 15610269
B	135	GLY	-	cloning artifact	GB 15610269
B	136	LEU	-	cloning artifact	GB 15610269
B	137	VAL	-	cloning artifact	GB 15610269
B	138	PRO	-	cloning artifact	GB 15610269
B	139	ARG	-	cloning artifact	GB 15610269
B	140	GLY	-	cloning artifact	GB 15610269
B	141	SER	-	cloning artifact	GB 15610269
B	142	HIS	-	cloning artifact	GB 15610269
B	143	MET	-	cloning artifact	GB 15610269

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	142.40Å 58.79Å 82.93Å 90.00° 125.50° 90.00°	Depositor
Resolution (Å)	50.00 – 3.10 50.00 – 3.10	Depositor EDS
% Data completeness (in resolution range)	79.7 (50.00-3.10) 79.7 (50.00-3.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 3.11Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.272 , 0.288 0.259 , 0.292	Depositor DCC
R_{free} test set	407 reflections (3.95%)	wwPDB-VP
Wilson B-factor (Å ²)	74.4	Xtriage
Anisotropy	0.262	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 95.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.448 for -h-2*1,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2018	wwPDB-VP
Average B, all atoms (Å ²)	89.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	C	0.30	0/551	0.78	0/848
2	D	0.27	0/579	0.79	0/892
3	A	0.57	0/508	0.86	0/681
3	B	0.54	0/508	0.84	0/681
All	All	0.44	0/2146	0.82	0/3102

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	492	0	271	18	0
2	D	516	0	282	14	0
3	A	505	0	530	23	0
3	B	505	0	530	23	0
All	All	2018	0	1613	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:21:DA:H2''	1:C:22:DA:H5'	1.42	1.01
3:B:160:LEU:HD12	3:B:170:ILE:HG23	1.66	0.76
3:A:202:VAL:HG13	3:B:205:THR:OG1	1.86	0.75
3:B:176:LEU:N	3:B:176:LEU:HD23	2.03	0.74
2:D:8:DG:H1'	2:D:9:DG:H5''	1.73	0.69
1:C:23:DG:OP1	3:A:197:ARG:HB3	1.95	0.66
2:D:29:DA:H2''	2:D:30:DA:C8	2.33	0.62
3:A:145:ASP:N	3:A:146:PRO:HD3	2.15	0.61
1:C:20:DA:C2	2:D:22:DA:C2	2.89	0.60
1:C:21:DA:H2''	1:C:22:DA:C5'	2.25	0.60
3:A:145:ASP:N	3:A:146:PRO:CD	2.64	0.60
3:B:145:ASP:N	3:B:146:PRO:HD3	2.16	0.60
3:A:151:THR:HG23	3:A:154:GLU:CD	2.27	0.60
2:D:29:DA:H2''	2:D:30:DA:H8	1.67	0.59
1:C:29:DA:H2''	1:C:30:DA:C8	2.38	0.58
3:B:145:ASP:N	3:B:146:PRO:CD	2.67	0.58
3:A:171:ALA:HB2	3:A:181:VAL:HG21	1.87	0.57
3:B:147:LEU:HB3	3:B:155:ARG:HD3	1.87	0.56
3:A:167:ASN:OD1	3:A:197:ARG:NH2	2.39	0.56
3:B:174:MET:O	3:B:175:PHE:HB2	2.06	0.56
1:C:10:DT:H2''	1:C:11:DT:H5''	1.89	0.55
1:C:29:DA:H2''	1:C:30:DA:H8	1.72	0.54
1:C:22:DA:H2''	1:C:23:DG:H5'	1.89	0.54
2:D:19:DT:H1'	2:D:20:DT:H5''	1.90	0.54
3:A:158:LEU:HG	3:A:204:ALA:HB2	1.90	0.53
2:D:21:DT:H1'	2:D:22:DA:H5'	1.92	0.52
3:B:158:LEU:HG	3:B:204:ALA:HB2	1.92	0.52
3:B:175:PHE:C	3:B:176:LEU:HD23	2.34	0.51
1:C:10:DT:H2''	1:C:11:DT:C6	2.45	0.51
2:D:21:DT:H2''	2:D:22:DA:H5'	1.92	0.51
3:A:166:THR:O	3:A:167:ASN:C	2.54	0.50
3:A:180:THR:O	3:A:184:TYR:HD2	1.96	0.49
3:A:176:LEU:N	3:A:176:LEU:HD23	2.28	0.49
3:A:196:ARG:HD2	3:B:166:THR:HG22	1.93	0.48
3:A:174:MET:O	3:A:175:PHE:HB2	2.13	0.48
2:D:21:DT:H1'	2:D:22:DA:C5'	2.44	0.47
3:B:156:THR:O	3:B:156:THR:HG22	2.14	0.46
3:A:198:THR:O	3:A:202:VAL:HG23	2.16	0.46
3:B:181:VAL:O	3:B:182:LYS:C	2.59	0.46
1:C:15:DG:H2''	1:C:16:DG:H5'	1.97	0.46
2:D:23:DG:OP1	3:B:197:ARG:HB3	2.16	0.46
1:C:19:DT:O5'	1:C:19:DT:H2'	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:147:LEU:HB3	3:A:155:ARG:HD3	1.99	0.45
3:A:160:LEU:HB3	3:A:170:ILE:HG12	1.98	0.45
3:A:189:LEU:HD21	3:A:197:ARG:HA	1.99	0.44
3:B:157:LEU:O	3:B:161:LEU:HB2	2.18	0.44
3:B:150:LEU:HD12	3:B:150:LEU:N	2.33	0.44
3:B:179:LYS:NZ	3:B:183:ASN:HD21	2.16	0.44
2:D:22:DA:H2''	2:D:23:DG:H5'	1.98	0.44
3:A:156:THR:HG22	3:A:156:THR:O	2.18	0.44
1:C:17:DA:N6	3:B:179:LYS:HE3	2.32	0.44
2:D:19:DT:H1'	2:D:20:DT:C5'	2.47	0.44
1:C:22:DA:C6	1:C:23:DG:C6	3.06	0.43
2:D:18:DC:H2''	2:D:19:DT:C6	2.54	0.43
3:A:166:THR:HG22	3:B:196:ARG:HD2	2.00	0.42
2:D:21:DT:C2'	2:D:22:DA:H5'	2.50	0.42
1:C:17:DA:H2''	1:C:18:DC:H5''	2.02	0.42
2:D:22:DA:C6	2:D:23:DG:C6	3.08	0.41
3:B:166:THR:O	3:B:167:ASN:C	2.64	0.41
3:B:179:LYS:HA	3:B:182:LYS:HD3	2.01	0.41
3:B:159:GLY:O	3:B:163:GLU:HG2	2.21	0.41
3:A:157:LEU:O	3:A:161:LEU:HB2	2.21	0.41
3:A:177:ALA:O	3:A:181:VAL:HG23	2.21	0.41
1:C:23:DG:H1'	1:C:24:DT:H5'	2.03	0.41
3:B:160:LEU:N	3:B:160:LEU:HD23	2.36	0.41
3:A:163:GLU:OE2	3:A:163:GLU:HA	2.21	0.40
1:C:28:DT:H2''	1:C:29:DA:H5''	2.02	0.40
3:B:179:LYS:HE2	3:B:182:LYS:HD3	2.03	0.40
1:C:22:DA:H2''	1:C:23:DG:C5'	2.51	0.40
1:C:23:DG:OP1	3:A:197:ARG:CB	2.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	63/95 (66%)	58 (92%)	5 (8%)	0	100	100
3	B	63/95 (66%)	60 (95%)	3 (5%)	0	100	100
All	All	126/190 (66%)	118 (94%)	8 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	54/80 (68%)	46 (85%)	8 (15%)	3	13
3	B	54/80 (68%)	50 (93%)	4 (7%)	13	40
All	All	108/160 (68%)	96 (89%)	12 (11%)	6	24

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	151	THR
3	A	153	GLN
3	A	161	LEU
3	A	172	ASP
3	A	176	LEU
3	A	196	ARG
3	A	197	ARG
3	A	208	LYS
3	B	160	LEU
3	B	176	LEU
3	B	179	LYS
3	B	196	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	153	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	A	183	ASN
3	B	183	ASN
3	B	199	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	24/43 (55%)	-1.43	0 100 100	47, 110, 150, 165	0
2	D	25/43 (58%)	-1.43	0 100 100	61, 109, 150, 172	0
3	A	65/95 (68%)	-1.52	0 100 100	24, 65, 103, 129	0
3	B	65/95 (68%)	-1.48	0 100 100	25, 61, 115, 168	0
All	All	179/276 (64%)	-1.48	0 100 100	24, 72, 144, 172	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.