



wwPDB EM Validation Summary Report ⓘ

Sep 9, 2026 – 04:53 PM EDT

PDB ID : 9ZP0 / pdb_00009zp0
EMDB ID : EMD-74512
Title : CryoEM structure of DNA-PK bound to N0 nucleosome
Authors : Lu, W.; He, Y.
Deposited on : 2025-12-16
Resolution : 4.11 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	5659 (3.62 - 4.61)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	I	147	
2	A	99	
2	E	99	
3	B	87	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	87	
4	C	111	
4	G	111	
5	D	96	
5	H	96	
6	J	147	
7	M	4128	
8	N	20	
9	K	609	
10	L	732	

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 97746 atoms, of which 47931 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (147-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
1	I	147	Total	C	H	N	O	P	0	0
			4682	1435	1651	566	883	147		

- Molecule 2 is a protein called Histone H3.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	A	99	Total	C	H	N	O	S	0	0
			1678	515	861	158	141	3		
2	E	98	Total	C	H	N	O	S	0	0
			1667	512	856	157	139	3		

- Molecule 3 is a protein called Histone H4.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	B	86	Total	C	H	N	O	S	0	0
			1453	440	755	141	116	1		
3	F	83	Total	C	H	N	O	S	0	0
			1372	418	710	129	114	1		

- Molecule 4 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	C	108	Total	C	H	N	O		0	0
			1701	520	876	159	146			
4	G	111	Total	C	H	N	O		0	0
			1783	539	925	169	150			

- Molecule 5 is a protein called Histone H2B.

Mol	Chain	Residues	Atoms						AltConf	Trace
5	D	96	Total	C	H	N	O	S	0	0
			1547	475	789	140	141	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						AltConf	Trace
5	H	95	Total	C	H	N	O	S	0	0
			1521	469	774	136	140	2		

- Molecule 6 is a DNA chain called DNA (147-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
6	J	147	Total	C	H	N	O	P	0	0
			4646	1423	1650	545	881	147		

- Molecule 7 is a protein called DNA-dependent protein kinase catalytic subunit.

Mol	Chain	Residues	Atoms						AltConf	Trace
7	M	3662	Total	C	H	N	O	S	0	0
			58977	18776	29693	4946	5370	192		

- Molecule 8 is a protein called Unknown peptide.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	N	20	Total	C	H	N	O	0	0
			141	60	40	20	21		

- Molecule 9 is a protein called X-ray repair cross-complementing protein 6.

Mol	Chain	Residues	Atoms						AltConf	Trace
9	K	490	Total	C	H	N	O	S	0	0
			7999	2533	4045	671	733	17		


- Molecule 10 is a protein called DNA repair protein Ku80.

Mol	Chain	Residues	Atoms						AltConf	Trace
10	L	535	Total	C	H	N	O	S	0	0
			8579	2736	4306	713	800	24		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (147-MER)



- Molecule 2: Histone H3




- Molecule 2: Histone H3



- Molecule 3: Histone H4








- Molecule 3: Histone H4





- Molecule 4: Histone H2A




- Molecule 4: Histone H2A




- Molecule 5: Histone H2B




- Molecule 5: Histone H2B




- Molecule 6: DNA (147-MER)






• Molecule 7: DNA-dependent protein kinase catalytic subunit







- Molecule 8: Unknown peptide

Chain N: 65% 35%

ASP	GLY	ILE	THR	LEU	ILE	THR	LYS	GLU	GLU	ALA	SER	GLY	SER	SER	VAL	THR	ALA	GLU	GLU	ALA	LYS	LYS	PHE	LEU	ALA	PRO	LYS	LYS	ASP	LYS	PRO	SER	GLY	ASP	THR	ALA	ALA	VAL	PHE	GLU	GLU	GLY	GLY	D724	V725	L728	L729	D730	M731	L732
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	96187	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	2000	Depositor
Maximum defocus (nm)	4000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	2.843	Depositor
Minimum map value	-0.364	Depositor
Average map value	0.008	Depositor
Map value standard deviation	0.063	Depositor
Recommended contour level	0.25	Depositor
Map size (Å)	446.4, 446.4, 446.4	wwPDB
Map dimensions	240, 240, 240	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.86, 1.86, 1.86	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	I	0.23	0/3403	0.47	0/5255
2	A	0.13	0/829	0.37	0/1111
2	E	0.17	0/823	0.47	0/1104
3	B	0.14	0/706	0.47	0/943
3	F	0.15	0/669	0.45	0/894
4	C	0.16	0/835	0.47	0/1127
4	G	0.14	0/868	0.40	0/1169
5	D	0.13	0/769	0.39	0/1032
5	H	0.13	0/758	0.33	0/1018
6	J	0.24	0/3357	0.47	0/5174
7	M	0.19	0/29884	0.48	3/40387 (0.0%)
9	K	0.15	0/4031	0.43	0/5429
10	L	0.16	0/4359	0.45	2/5881 (0.0%)
All	All	0.19	0/51291	0.47	5/70524 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	E	0	1
3	B	0	1
4	C	0	1
7	M	0	8
9	K	0	1
All	All	0	12

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	L	339	CYS	CA-C-N	6.74	133.82	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	L	339	CYS	C-N-CA	6.74	133.82	121.70
7	M	1221	ILE	N-CA-C	-6.34	106.79	112.12
7	M	190	ILE	N-CA-C	-6.24	105.26	111.88
7	M	212	VAL	N-CA-C	-5.80	107.03	111.62

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	B	19	ARG	Sidechain
4	C	42	ARG	Peptide
2	E	38	PRO	Peptide
7	M	185	HIS	Peptide
7	M	2197	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	I	3031	1651	1651	96	0
2	A	817	861	858	26	0
2	E	811	856	853	25	0
3	B	698	755	752	30	0
3	F	662	710	709	26	0
4	C	825	876	875	30	0
4	G	858	925	922	13	0
5	D	758	789	786	22	0
5	H	747	774	773	16	0
6	J	2996	1650	1650	102	0
7	M	29284	29693	29680	873	0
8	N	101	40	24	4	0
9	K	3954	4045	4042	142	0
10	L	4273	4306	4303	128	0
All	All	49815	47931	47878	1388	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 1388 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:M:2216:LEU:HD21	7:M:2241:LEU:HD22	1.45	0.96
4:C:17:ARG:NH1	6:J:45:DC:OP1	2.02	0.92
7:M:3499:ILE:HD11	7:M:3521:ILE:HD13	1.52	0.91
7:M:879:MET:HE3	7:M:879:MET:O	1.71	0.90
7:M:2424:MET:O	7:M:2432:GLN:NE2	2.04	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	97/99 (98%)	89 (92%)	8 (8%)	0	100	100
2	E	96/99 (97%)	86 (90%)	9 (9%)	1 (1%)	12	46
3	B	84/87 (97%)	82 (98%)	1 (1%)	1 (1%)	10	42
3	F	81/87 (93%)	78 (96%)	3 (4%)	0	100	100
4	C	106/111 (96%)	86 (81%)	19 (18%)	1 (1%)	14	48
4	G	109/111 (98%)	99 (91%)	9 (8%)	1 (1%)	14	48
5	D	94/96 (98%)	87 (93%)	7 (7%)	0	100	100
5	H	93/96 (97%)	91 (98%)	2 (2%)	0	100	100
7	M	3632/4128 (88%)	3203 (88%)	412 (11%)	17 (0%)	24	61
9	K	486/609 (80%)	429 (88%)	54 (11%)	3 (1%)	21	58
10	L	529/732 (72%)	464 (88%)	61 (12%)	4 (1%)	16	52
All	All	5407/6255 (86%)	4794 (89%)	585 (11%)	28 (0%)	26	61

5 of 28 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	G	18	SER
7	M	186	PRO
7	M	189	MET
7	M	1479	VAL
7	M	1551	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	86/86 (100%)	86 (100%)	0	100	100
2	E	86/86 (100%)	86 (100%)	0	100	100
3	B	72/72 (100%)	72 (100%)	0	100	100
3	F	68/72 (94%)	68 (100%)	0	100	100
4	C	84/88 (96%)	84 (100%)	0	100	100
4	G	88/88 (100%)	88 (100%)	0	100	100
5	D	82/82 (100%)	82 (100%)	0	100	100
5	H	81/82 (99%)	81 (100%)	0	100	100
7	M	3266/3671 (89%)	3266 (100%)	0	100	100
9	K	444/548 (81%)	444 (100%)	0	100	100
10	L	481/649 (74%)	481 (100%)	0	100	100
All	All	4838/5524 (88%)	4838 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 46 such sidechains are listed below:

Mol	Chain	Res	Type
7	M	3133	GLN
7	M	3590	ASN
7	M	3154	GLN
7	M	3510	GLN
7	M	3760	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-74512. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 120



Y Index: 120




Z Index: 120

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 108



Y Index: 111




Z Index: 101


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.25. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 912 nm³; this corresponds to an approximate mass of 824 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.243 $\text{\AA}^{-1}$


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-74512 and PDB model 9ZP0. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.25 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.25).

9.4 Atom inclusion [i](#)



At the recommended contour level, 92% of all backbone atoms, 92% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.25) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9190	 0.2470
A	 0.7460	 0.0030
B	 0.7810	 0.0420
C	 0.7170	 0.0480
D	 0.8220	 0.0820
E	 0.7960	 0.0400
F	 0.8590	 0.0560
G	 0.7470	 0.0420
H	 0.8320	 0.0510
I	 0.8620	 0.1070
J	 0.8440	 0.0960
K	 0.9670	 0.2100
L	 0.9330	 0.1500
M	 0.9540	 0.3380
N	 0.9900	 0.4440

