



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:02 AM UTC

PDB ID : 4ZWP / pdb_00004zwp
Title : Crystal structure of organophosphate anhydrolase/prolidase mutant Y212F
Authors : Daczkowski, C.M.; Pegan, S.D.; Harvey, S.P.
Deposited on : 2015-05-19
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	77% 18% .
1	B	440	2% 82% 15% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7338 atoms, of which 8 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called OPAA organophosphate prolidase anhydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	422	Total	C	N	O	S	0	1	0
			3424	2195	583	633	13			
1	B	428	Total	C	N	O	S	0	4	0
			3491	2232	597	649	13			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	210	THR	ASN	variant	UNP Q44238
A	211	PRO	ALA	variant	UNP Q44238
A	212	PHE	TYR	engineered mutation	UNP Q44238
A	314	ASN	ASP	variant	UNP Q44238
A	438	GLU	-	expression tag	UNP Q44238
A	439	LEU	-	expression tag	UNP Q44238
A	440	ASP	-	expression tag	UNP Q44238
B	210	THR	ASN	variant	UNP Q44238
B	211	PRO	ALA	variant	UNP Q44238
B	212	PHE	TYR	engineered mutation	UNP Q44238
B	314	ASN	ASP	variant	UNP Q44238
B	438	GLU	-	expression tag	UNP Q44238
B	439	LEU	-	expression tag	UNP Q44238
B	440	ASP	-	expression tag	UNP Q44238


- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Mn	0	0
			4	4		
2	B	3	Total	Mn	0	0
			3	3		

- Molecule 3 is BARIUM ION (CCD ID: BA) (formula: Ba).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Ba	0	0
			3	3		

- Molecule 4 is N,N'-bis(1-methylethyl)phosphorodiamidic acid (CCD ID: M44) (formula: $C_6H_{17}N_2O_2P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			11	6	2	2	1		
4	B	1	Total	C	N	O	P	0	0
			11	6	2	2	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	182	Total	O	0	0
			182	182		
6	B	195	Total	O	0	0
			195	195		

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 

- Chain B:
-
- 2% 82% 15%
- Met N2 K3 L4 L15 T19 I23 E24 V32 Y47 L60 I63 I70 D87 F88 W89 H90 LYS VAL PRO ASP GLU P96 D102 V110 K111 P112 L119 P120 L133 M144 PHE E164 L165 M168 N172 F193 Q197 F212 T215 N219 E220 A223 H229 R239 S240 F241 I243 G246 Y252 A253 A254 T257 D261 G264 Q285 D298 I320 T326 P331 L340 F348 P359 P360 GLU GLY HIS PRO PHE LEU R367 T379 I380 E381 D393 L394 A395 N400

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.49Å 68.18Å 142.48Å 90.00° 110.50° 90.00°	Depositor
Resolution (Å)	48.94 – 2.40 48.94 – 2.40	Depositor EDS
% Data completeness (in resolution range)	94.7 (48.94-2.40) 92.3 (48.94-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 2.39Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.173 , 0.226 0.174 , 0.225	Depositor DCC
R_{free} test set	1990 reflections (5.67%)	wwPDB-VP
Wilson B-factor (Å ²)	32.7	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.025 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7338	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: M44, MN, BA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/3513	0.76	1/4763 (0.0%)
1	B	0.48	0/3590	0.77	0/4870
All	All	0.49	0/7103	0.77	1/9633 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	219	ASN	CB-CA-C	-5.55	109.68	117.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3424	0	3281	59	0
1	B	3491	0	3332	49	0
2	A	4	0	0	0	0
2	B	3	0	0	0	0
3	A	3	0	0	0	0
4	A	11	0	16	2	0
4	B	11	0	16	4	0
5	B	6	8	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	182	0	0	17	0
6	B	195	0	0	10	0
All	All	7330	8	6653	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:393:ASP:OD2	6:B:601:HOH:O	1.86	0.93
1:A:226:HIS:CD2	1:B:89:TRP:HB3	2.15	0.82
1:A:427:GLU:OE2	6:A:601:HOH:O	1.98	0.82
1:B:47:TYR:OH	6:B:602:HOH:O	1.98	0.79
1:B:331:PRO:O	6:B:603:HOH:O	2.01	0.78
1:A:86:VAL:O	6:A:602:HOH:O	2.04	0.76
1:B:298:ASP:OD1	6:B:604:HOH:O	2.03	0.75
1:B:212:PHE:CD2	4:B:504:M44:H5	2.22	0.75
1:B:285:GLN:NE2	6:B:606:HOH:O	2.19	0.75
1:A:212:PHE:CE2	4:A:508:M44:H3	2.21	0.74
1:A:387:ILE:HG21	1:B:88:PHE:HB2	1.72	0.72
1:A:137:GLN:HG2	1:A:144:MET:HE1	1.76	0.67
1:A:401:GLN:OE1	6:A:606:HOH:O	2.13	0.66
1:A:440:ASP:O	6:A:605:HOH:O	2.13	0.66
1:A:352:GLU:OE2	6:A:607:HOH:O	2.13	0.66
1:A:76:ASP:O	6:A:608:HOH:O	2.14	0.65
1:A:183:ARG:HG3	1:A:260:TYR:CE1	2.31	0.65
1:B:440:ASP:OXT	6:B:605:HOH:O	2.14	0.65
4:B:504:M44:HN2	4:B:504:M44:H3A	1.62	0.65
1:B:381:GLU:HB3	1:B:420:GLU:HB2	1.78	0.64
1:A:226:HIS:ND1	6:A:616:HOH:O	2.28	0.64
1:A:266:GLY:N	6:A:615:HOH:O	2.28	0.62
1:B:63:ILE:HG12	6:B:668:HOH:O	2.01	0.61
1:A:165:LEU:HD22	1:A:424:ILE:HD13	1.82	0.61
1:A:349:MET:HE3	1:A:354:GLY:HA2	1.83	0.60
1:A:125:ARG:N	6:A:604:HOH:O	2.12	0.60
1:A:183:ARG:HG3	1:A:260:TYR:CZ	2.38	0.58
1:B:229[B]:HIS:ND1	6:B:614:HOH:O	2.32	0.58
1:B:164:GLU:HG2	1:B:252:TYR:OH	2.04	0.56
1:B:404:ASN:O	1:B:408:VAL:HG23	2.05	0.56
1:B:23:ILE:HG13	1:B:24:GLU:N	2.20	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:193:PHE:O	1:B:197:GLN:HG2	2.06	0.56
1:A:373:GLU:OE1	6:A:610:HOH:O	2.18	0.55
1:B:164:GLU:HG2	1:B:252:TYR:CZ	2.41	0.55
1:A:21:GLU:OE2	1:A:25:ARG:NH2	2.41	0.54
1:B:19:THR:O	1:B:23:ILE:HG23	2.08	0.54
1:A:206:SER:HA	6:A:705:HOH:O	2.08	0.54
1:B:212:PHE:CE2	4:B:504:M44:H5	2.42	0.54
1:A:164:GLU:HG2	1:A:252:TYR:OH	2.08	0.53
1:A:221:ASN:HA	1:A:224:ILE:HD12	1.88	0.53
1:B:432:ASN:O	1:B:436:GLU:HG3	2.09	0.53
1:A:110:VAL:HG13	1:A:111:LYS:HG2	1.91	0.53
1:A:63:ILE:O	6:A:609:HOH:O	2.18	0.53
1:A:242:LEU:C	1:A:242:LEU:HD23	2.34	0.53
1:B:220:GLU:OE2	1:B:239:ARG:NH2	2.42	0.52
1:A:69:TRP:HB2	1:A:81:ILE:HB	1.91	0.52
1:B:395:ALA:HA	1:B:400:ASN:OD1	2.09	0.52
1:A:349:MET:HE3	1:A:354:GLY:CA	2.39	0.52
1:A:231:ASP:HA	6:A:642:HOH:O	2.10	0.52
1:A:256:ILE:HD13	1:A:434:THR:HG23	1.91	0.52
1:B:119:LEU:HB3	1:B:120:PRO:HD2	1.90	0.52
1:A:407:LYS:HE2	1:A:410:GLU:OE1	2.11	0.51
1:B:32:VAL:HA	1:B:70:ILE:O	2.11	0.50
1:A:291:LEU:HD23	1:A:371:LYS:HG3	1.93	0.50
1:A:136:ALA:CB	1:A:144:MET:HE2	2.41	0.49
1:B:60:LEU:HD12	1:B:60:LEU:N	2.27	0.49
1:B:438:GLU:HA	6:B:632:HOH:O	2.12	0.49
4:B:504:M44:H3A	4:B:504:M44:N2	2.26	0.49
1:A:212:PHE:CE1	4:A:508:M44:H2	2.48	0.48
1:B:241:PHE:CZ	1:B:243:ILE:HB	2.48	0.48
1:B:257:THR:HB	1:B:420:GLU:HB3	1.95	0.48
1:A:421:ASP:OD2	1:A:435:ARG:NE	2.47	0.48
1:A:193:PHE:O	1:A:197:GLN:HG2	2.13	0.48
1:B:102:ASP:OD1	1:B:102:ASP:N	2.40	0.48
1:A:147:GLU:HB3	1:A:148:PRO:HD3	1.96	0.47
1:B:164:GLU:O	1:B:168:MET:HG3	2.14	0.47
1:A:320:ILE:HG22	1:A:326:THR:HG23	1.97	0.47
1:B:380:ILE:O	1:B:420:GLU:HA	2.15	0.47
1:A:299:CYS:O	1:A:303:VAL:HG23	2.15	0.46
1:B:261:ASP:OD1	1:B:264:GLY:N	2.41	0.46
1:B:133:LEU:HD22	1:B:144:MET:HB3	1.98	0.46
1:A:136:ALA:HB3	1:A:144:MET:HE2	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:GLN:NE2	1:A:78:PRO:HD3	2.31	0.46
1:A:320:ILE:HG22	1:A:326:THR:CG2	2.46	0.46
1:A:423:ILE:HD12	1:A:423:ILE:N	2.31	0.45
1:A:168:MET:HE1	1:A:338:ILE:HG23	1.98	0.45
1:A:381:GLU:HB3	1:A:420:GLU:HB2	1.98	0.45
1:B:165:LEU:HD22	1:B:424:ILE:HD13	1.98	0.45
1:A:147:GLU:HG2	1:A:151:ASN:ND2	2.32	0.45
1:A:352:GLU:HG3	6:A:716:HOH:O	2.16	0.45
1:A:366:LEU:HD23	1:A:369:THR:HG22	1.99	0.44
1:A:229[A]:HIS:CG	1:A:230:PHE:N	2.85	0.43
1:B:119:LEU:HB3	1:B:120:PRO:CD	2.48	0.43
1:A:215:ILE:HB	1:A:244:ASP:HB3	2.00	0.43
1:A:134:GLU:N	1:A:134:GLU:OE1	2.51	0.43
1:B:246:GLY:HA3	1:B:254:ALA:O	2.19	0.43
1:A:401:GLN:HG2	6:A:603:HOH:O	2.17	0.43
1:A:349:MET:HE2	1:A:351:ASP:O	2.17	0.43
1:B:87:ASP:OD1	1:B:89:TRP:HB2	2.18	0.43
1:B:320:ILE:HG22	1:B:326:THR:HG22	2.01	0.43
1:A:26:GLU:OE1	6:A:611:HOH:O	2.21	0.42
1:A:190:LYS:HB3	1:A:194:GLU:HB2	2.01	0.42
1:B:412:LYS:N	1:B:413:PRO:CD	2.83	0.42
1:B:427:GLU:HG3	1:B:428[B]:ASP:N	2.35	0.42
1:B:215:ILE:O	1:B:243:ILE:HA	2.20	0.42
1:A:246:GLY:HA3	1:A:254:ALA:O	2.20	0.42
1:B:367:ARG:HE	1:B:367:ARG:HB2	1.69	0.42
1:A:28:LEU:HA	1:A:125:ARG:HG2	2.01	0.42
1:B:223:ALA:HB2	1:B:415:GLY:O	2.19	0.42
1:B:15:LEU:HD23	1:B:15:LEU:HA	1.83	0.41
1:B:379:THR:HA	1:B:421:ASP:O	2.20	0.41
1:B:219:ASN:HA	1:B:240:SER:OG	2.21	0.41
1:B:367:ARG:HB3	6:B:634:HOH:O	2.21	0.41
1:A:9:ALA:HB2	1:A:103:TYR:OH	2.21	0.41
1:A:243:ILE:O	1:A:257:THR:HA	2.20	0.41
1:B:111:LYS:HA	1:B:112:PRO:HD3	1.89	0.40
1:B:252:TYR:CE2	1:B:340:LEU:HG	2.56	0.40
1:A:183:ARG:NH2	6:A:631:HOH:O	2.50	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	417/440 (95%)	410 (98%)	7 (2%)	0	100	100
1	B	426/440 (97%)	418 (98%)	8 (2%)	0	100	100
All	All	843/880 (96%)	828 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/373 (96%)	352 (99%)	5 (1%)	59	79
1	B	366/373 (98%)	363 (99%)	3 (1%)	73	86
All	All	723/746 (97%)	715 (99%)	8 (1%)	65	82

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ILE
1	A	97	ASN
1	A	125	ARG
1	A	172	ASN
1	A	327	SER
1	B	4	LEU
1	B	60	LEU
1	B	172	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	GLN
1	A	177	GLN
1	A	208	ASN
1	B	55	GLN
1	B	204	GLN
1	B	208	ASN
1	B	214	ASN
1	B	285	GLN
1	B	296	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 10 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	B	505	-	5,5,5	0.49	0	5,5,5	0.76	0
4	M44	B	504	2	7,10,10	3.11	1 (14%)	7,14,14	0.49	0
4	M44	A	508	2	7,10,10	3.16	1 (14%)	7,14,14	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	505	-	-	0/4/4/4	-
4	M44	B	504	2	-	4/8/10/10	-
4	M44	A	508	2	-	0/8/10/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	508	M44	P-O2	8.21	1.60	1.46
4	B	504	M44	P-O2	8.08	1.60	1.46

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	504	M44	C4-N2-P-N1
4	B	504	M44	C5-C4-N2-P
4	B	504	M44	C6-C4-N2-P
4	B	504	M44	C1-N1-P-O2

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	504	M44	4	0
4	A	508	M44	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	422/440 (95%)	-0.07	4 (0%) 81 78	19, 36, 61, 76	1 (0%)
1	B	428/440 (97%)	-0.06	8 (1%) 66 62	20, 34, 64, 87	4 (0%)
All	All	850/880 (96%)	-0.07	12 (1%) 73 69	19, 36, 63, 87	5 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	365	PHE	3.3
1	A	89	TRP	3.1
1	A	313	VAL	2.9
1	B	96	PRO	2.8
1	B	359	PRO	2.8
1	A	355	ALA	2.7
1	B	348	PHE	2.6
1	B	360	PRO	2.4
1	B	102	ASP	2.1
1	B	229[A]	HIS	2.1
1	B	367	ARG	2.1
1	B	110	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	B	505	6/6	0.77	0.17	45,54,57,58	0
3	BA	A	503	1/1	0.78	0.09	163,163,163,163	1
3	BA	A	507	1/1	0.89	0.08	156,156,156,156	0
3	BA	A	506	1/1	0.92	0.08	110,110,110,110	0
4	M44	A	508	11/11	0.93	0.14	38,43,80,81	0
4	M44	B	504	11/11	0.94	0.14	31,53,75,79	0
2	MN	A	504	1/1	0.97	0.04	45,45,45,45	0
2	MN	A	501	1/1	0.99	0.04	27,27,27,27	0
2	MN	A	505	1/1	0.99	0.03	34,34,34,34	0
2	MN	B	502	1/1	0.99	0.03	28,28,28,28	0
2	MN	B	503	1/1	0.99	0.06	46,46,46,46	0
2	MN	A	502	1/1	0.99	0.04	29,29,29,29	0
2	MN	B	501	1/1	1.00	0.03	26,26,26,26	0

6.5 Other polymers

There are no such residues in this entry.